

UNIVERSIDADE DE LISBOA
FACULDADE DE CIÊNCIAS
DEPARTAMENTO DE BIOLOGIA VEGETAL



Sensitivity profile of triple-negative breast cancer cell lines to combination therapies including RANKL blockade

João Brito Colaço Carvalho Belo

Mestrado em Biologia Molecular e Genética

Dissertação orientada por:

Professora Doutora Sandra Cristina Cara de Anjo Casimiro, IMM-FMUL

Professora Doutora Maria Helena de Figueiredo Ramos Caria, FCUL

2020

ACKNOWLEDGEMENTS

I would like to start off by thanking everyone that, one way or another, helped me get through this year successfully. Turns out, developing a research project was very different from what I imagined. It really did “take a village”! Here I would like to personally thank some of the people who were absolutely crucial in this special and challenging, but also overall fun, time.

To Professor Luís Costa, thank you for accepting me in your amazing lab and giving me the opportunity to meet an incredible group of people who are and will, no doubt, leave their mark and change this world for the better – with you “on the wheel”. It was very inspiring to watch you discussing experimental data and figuring out ways to improve the therapeutic options of your (and all) cancer patients.

To Professor Sandra Casimiro, who so kindly gave me this project and guided me through every step of it. Thank you for your endless patience, for your dedication in making me both a better professional and a better person. For all the mistakes I made, and also the ones I didn’t, I am deeply grateful for all the time you spent helping me reach this new milestone. I know I had a lot of “rough edges” when I first came to the lab, and I will certainly leave with some, but I feel like I end this chapter of my life a lot better than I started it. And that was in no small part thanks to you.

To Professor Helena Caria, thank you for accepting to be my internal supervisor and for all the availability and kindness throughout this year.

I would also like to thank Inês. Without you, I couldn’t have done this. Really. Thank you for all the hours you spent teaching me everything I needed to know. For helping me feel comfortable in a space that was quite intimidating at first. For always encouraging me and helping me correct my mistakes and learn from them. And most of all, for simply being there, and making sure I knew I could count on you for even the smallest thing.

To everyone from the LCosta Lab. Raquel, Patrícias, Marta, Ana, Inês, Sandra (yes, again!). I know I’m not the most sociable person. But from the simple good mornings with a smile, the lunches and snacks in the cafeteria, the “is everything ok?”s, the conversations in the lab while refilling tips, the availability to help me at any time, and so much more; everything made me feel welcome and it really did make a difference. There is not a single person from the lab I cannot imagine a great future for. Everyone just seems so effortlessly competent and intelligent, I really enjoyed just listening to you discussing science. I have no doubts you’ll continue on having the interesting careers I caught a glimpse of during this year. And I wish you all the luck in that, and hope our paths cross again in some way.

And finally, last but not least, I’d like to thank my friends and family (and dog!). I put you all here together not only because it would be a bit lame to start reciting names, but also because I’m lucky enough to be surrounded with friends who are like family and family who are friends. As long as I’ve got your support, I can do everything I set my mind to. Thank you for putting up with me every day and for sharing the good and bad moments of my life. For the love and happiness you give me, and for believing in me no matter what.

Thank you.

This project was funded by the research project PTDC/MED-ONC/28636/2017 from Fundação para a Ciência e a Tecnologia (FCT)/Ministério da Ciência, Tecnologia e Ensino Superior (MCTES) through Fundos do Orçamento de Estado.

RESUMO

O cancro da mama (CM) é o cancro com a segunda mais elevada mortalidade a nível mundial. Dada a heterogeneidade que apresenta, é classificado em subtipos de acordo com a expressão de determinados recetores moleculares, de modo a permitir uma maior especificidade, e consequentemente eficácia, no tratamento dos pacientes. Um dos subtipos definidos é o CM triplo negativo (TNBC, do inglês *triple-negative breast cancer*), caracterizado pela ausência de expressão de recetores de progesterona (RP) e de recetores de estrogénio (RE), e ainda por não apresentar amplificação do recetor 2 do fator de crescimento epidérmico humano (HER2).

O TNBC constitui cerca de 10% a 15% dos casos de CM e está associado a um prognóstico pior do que os restantes subtipos, possuindo os pacientes taxas de sobrevivência inferiores e maior probabilidade de recidiva. Apesar de o TNBC apresentar uma elevada sensibilidade à quimioterapia, sendo esta a terapia padrão para os doentes com TNBC, a aquisição de mecanismos de resistência explica as recidivas e pior prognóstico observados. Deste modo, é necessária a investigação de novas abordagens clínicas direcionadas que reduzam o impacto negativo do TNBC na sociedade.

Os inibidores da poli-(ADP-ribose) polimerase (PARP) (iPARP) são uma classe de fármacos que afeta a reparação de quebras simples da cadeia de DNA (SSB, do inglês *single strand breaks*). Em paralelo, CM com mutações nos genes *BRCA1* e/ou *BRCA2* tendem a perder o alelo saudável, o que lhes confere deficiência na reparação por recombinação homóloga (HRR, do inglês *homologous recombination repair*), impedindo a correção de quebras duplas da cadeia de DNA (DSB, do inglês *double strand breaks*). Estes dois défices resultam eventualmente em apoptose. O olaparib é um iPARP aprovado para tratamento de CM, nomeadamente em doentes com CM HER2⁻ metastático ou avançado com mutações germinais em *BRCA1/2* e de doentes com CM com recetores hormonais (RH) não responsivos a terapia hormonal (TH). É possível que outros doentes com TNBC beneficiem do olaparib, particularmente com tumores possuindo mutações somáticas em *BRCA1/2* ou metilação do promotor destes genes.

Por outro lado, os inibidores da fosfoinositídeo 3-quinase (PI3K) (iPI3K) são fármacos que atuam ao nível da via do PI3K/Akt/mTOR, que é uma via de sinalização frequentemente mutada em CM, incluindo TNBC. Enquanto que o PI3K promove a ativação da quinase Akt, a fosfatase homóloga à tensina (PTEN) regula negativamente este processo, estando associada a funções de supressão tumoral. Recentemente, foi aprovado um iPI3K, o alpelisib, para tratamento de doentes com CM RE⁺, RP⁺, HER2⁻ metastático ou avançado com mutações no gene *PIK3CA* e após progressão sob TH. Este inibidor apresenta especificidade para a isoforma α do PI3K, cuja subunidade catalítica é codificada pelo gene *PIK3CA*, e atua através de um mecanismo pró-apoptótico. É possível que o alpelisib tenha utilidade no tratamento de doentes com TNBC com mutações ao nível do *PIK3CA*.

Curiosamente, foi previamente descrita uma sinergia entre iPARP e iPI3K em modelos de TNBC, o que levou a um ensaio clínico que pretendeu testar a combinação de olaparib com os iPI3K buparlisib ou alpelisib em doentes com TNBC recorrente ou cancro ovárico seroso de alto grau. Este ensaio demonstrou existir sinergia entre olaparib e alpelisib em cancro ovárico HRR-competente. O mesmo poderá ser verdade para TNBC *BRCA*-competente, e talvez especialmente em doentes com mutações no *PIK3CA*.

O ligando do recetor ativador do fator nuclear kappa B (RANKL) é uma proteína homotrimérica pertencente à superfamília dos fatores de necrose tumoral (TNF, do inglês *tumour necrosis factors*) e que pode existir na forma membranar ou solúvel. O RANKL liga-se ao recetor RANK, um recetor homotrimérico transmembranar que requer proteínas adaptadoras para ser funcional. Destas, destaca-se o fator associado a TNF 6 (TRAF6), que tem um envolvimento crucial na ativação de vias a jusante de

RANKL/RANK, nomeadamente as vias do PI3K/Akt/mTOR, das proteínas quinases ativadas por mitogénios (MAPKs), e da Src/fosfolipase gamma (PLC γ). Esta cascata de transdução de sinal resulta na promoção da sobrevivência, diferenciação e proliferação celulares devido à translocação nuclear de ativadores transcricionais, nomeadamente o NF- κ B. A via de sinalização RANKL/RANK é negativamente regulada pela osteoprotegerina (OPG), um recetor “engodo” que é capaz de se ligar ao RANKL com uma afinidade largamente superior à que o RANK apresenta.

A via do RANKL/RANK está envolvida em diversos processos fisiológicos, sendo de ressaltar a importância crucial na manutenção da homeostase óssea. Uma outra função crítica desta via ocorre ao nível do tecido mamário, nomeadamente promovendo a morfogénese da glândula mamária durante a gravidez. A progesterona possui um efeito mitogénico nas células epiteliais mamárias, ligando-se ao RP de células epiteliais luminais e promovendo um aumento da expressão de RANKL. Esta citocina vai interagir com células RANK⁺, tanto basais como luminais, e promover fenómenos de proliferação e sobrevivência. De destaque, há duas cascatas de sinalização envolvidas neste processo, uma resultando na transcrição da ciclina D1 dependente do NF- κ B, e outra resultando na redução da transcrição da p21 mediada pelo inibidor da proteína *DNA binding 2* (Id2).

No contexto oncológico, já foi observada expressão de RANK e RANKL em diversos tumores humanos e, inclusive, foram propostas associações preditivas de mau prognóstico. O envolvimento da via RANKL/RANK neste contexto foi já descrito ao nível da promoção da transição epitélio-mesénquima (TEM), migração celular, neovascularização tumoral, estabelecimento de metástases à distância, imunossupressão, entre outros. Um dos cancros em que foi estabelecida uma relação entre esta via e a carcinogénese é o CM – tendo sido destacado o envolvimento da progesterona e derivados sintéticos. Para além disso, foi ainda identificado o papel crucial do eixo progesterona/RANKL na tumorigénese de CM com mutações em *BRCA1*, que são maioritariamente TNBC. Ainda no contexto do CM, a via RANKL/RANK está por detrás do ciclo vicioso que se observa frequentemente nas metástases ósseas, em que ocorre uma promoção descontrolada da reabsorção óssea, criando no osso um nicho favorável ao estabelecimento das células tumorais.

A via RANKL/RANK é, deste modo, um alvo terapêutico emergente tanto ao nível do TNBC como do CM no geral. Entre 2010 e 2011, foi aprovado para utilização clínica um anticorpo monoclonal anti-RANKL totalmente humanizado, denosumab, sendo atualmente utilizado em doentes com CM sob terapia de ablação hormonal e em contexto de doença metastática óssea. Tendo sido previamente observado um efeito do bloqueio do RANKL *in vitro* na prevenção da tumorigénese de CM mutado em *BRCA1*, é possível que este bloqueio se possa demonstrar vantajoso em TNBC quando integrado em abordagens terapêuticas combinatórias.

Neste projeto, colocámos a hipótese de que a inibição farmacológica da via RANKL/RANK, através do bloqueio do RANKL, poderia apresentar sinergia com iPARP, como o olaparib, e/ou iPI3K, como o alpelisib, no contexto do TNBC.

Considerando que o TNBC pode expressar RANK e RANKL, e que uma maior expressão destas proteínas poderá estar associada a uma maior suscetibilidade ao bloqueio por RANKL, começámos por avaliar a expressão de ambas num painel de linhas celulares TNBC (BT-20, HCC1937, MDA-MB-231 e MDA-MB-468).

De seguida, de modo a justificar uma abordagem de inibição da via RANKL/RANK, foi confirmada a ativação da via por RANKL exógeno, nomeadamente a fosforilação de proteínas a jusante da via como p-65, Erk e Akt. Para realizar o bloqueio do RANKL, foi selecionada a proteína recombinante OPG-Fc, que confirmámos ser capaz de inibir a ativação da via pelo RANKL.

Estando verificada a possibilidade da abordagem inibitória, foram primeiramente realizados ensaios de viabilidade celular após 72 horas de exposição a olaparib, alpelisib e OPG-Fc, em regime individual. As diferentes linhas celulares mostraram-se, na sua maioria, resistentes ao olaparib, mas sensíveis ao tratamento com alpelisib, embora a diferentes níveis. A OPG-Fc não mostrou afetar, *per se*, a viabilidade celular. Finalmente, de modo a verificar o efeito do bloqueio do RANKL quando em combinação com olaparib e/ou alpelisib, foram realizados ensaios de viabilidade envolvendo combinações duplas de olaparib e OPG-Fc ou alpelisib e OPG-Fc; e combinações triplas de olaparib, alpelisib e OPG-Fc. Nenhuma das referidas combinações se mostrou significativamente vantajosa em termos de redução da viabilidade celular quando comparados os regimes com e sem OPG-Fc. Foi confirmado por análise de expressão proteica que o olaparib e o alpelisib estavam, de facto a afetar os respetivos alvos moleculares, confirmando-se assim que a adição de OPG-Fc não foi eficaz no contexto das células TNBC avaliadas.

Apesar dos resultados obtidos, a inibição da via do RANKL/RANK continua a ser um alvo terapêutico importante no âmbito do TNBC, existindo evidências pré-clínicas de sinergia do bloqueio do RANKL com inibidores de *checkpoints* do sistema imunitário.

Palavras-chave: cancro da mama triplo negativo; via RANKL/RANK; olaparib; alpelisib; OPG-Fc.

ABSTRACT

Triple-negative breast cancer (TNBC) accounts for around 10% to 15% of breast cancer (BC) cases and is typically associated with bad clinical outcomes. Chemotherapy is the current standard of care for this complex disease. However, high likelihood of relapse is observed due to acquired chemoresistance. Moreover, TNBC also strikingly lacks effective targeted therapies. These aspects highlight the critical importance of research on new and innovative approaches to treat these patients. Combination therapies are quite pertinent for drug research on clinically relevant alternatives. Namely, they may allow drug repurposing for faster yet effective translation into the clinical practice and also permit the administration of lower drug dosages, thus reducing side effects.

The receptor activator of nuclear factor kappa B ligand (RANKL)/RANK pathway became an emerging target in this context after being linked to TNBC tumorigenesis. This pathway is involved in cell proliferation and survival mainly through downstream activation of the NF- κ B, phosphoinositide 3-kinase (PI3K)/Akt/mammalian target of rapamycin (mTOR) and mitogen-activated protein kinase (MAPK) pathways. Furthermore, dual RANKL and RANK expression was associated with worse prognosis amongst TNBC patients when compared to TNBC presenting RANK single expression. Here, we describe the impact of a combination regimen including RANKL blockade with osteoprotegerin (OPG)-Fc and poly (adenosine diphosphate-ribose) polymerase (PARP) inhibition with olaparib and/or PI3K inhibition with alpelisib in TNBC cell lines BT-20, HCC1937, MDA-MB-231 and MDA-MB-468. The cell line panel was initially characterised according to RANK and RANKL expression. Following exogenous RANKL stimulus, we verified pathway activation, which was successfully abrogated upon OPG-Fc treatment. We report no statistically significant impact on cell viability associated with combination therapies including RANKL blockade, PARP inhibition and/or PI3K inhibition *in vitro* in the TNBC cell lines.

Nonetheless, despite the lack of synergy observed in the selected combinations, RANK signalling remains an important target in TNBC patients, with reports of pre-clinical success with other drug types, namely immune checkpoint inhibitors.

Keywords: triple-negative breast cancer; RANKL/RANK pathway; olaparib; alpelisib; OPG-Fc.

INDEX

ACKNOWLEDGEMENTS	ii
RESUMO	iii
ABSTRACT	vi
INDEX	vii
LIST OF FIGURES	viii
LIST OF TABLES	ix
LIST OF ABBREVIATIONS	x
1. STATE OF THE ART.....	1
1.1. Triple-negative breast cancer	1
1.1.1. PARP inhibitors (PARPi) in the treatment of TNBC	2
1.1.2. PI3K inhibitors (PI3Ki) in the treatment of TNBC	2
1.1.3. PARPi and PI3Ki synergy	3
1.2. RANKL/RANK pathway	3
1.2.1. Physiological functions	4
1.2.2. Involvement in cancer	6
1.2.3. Pathway blockade.....	8
2. AIMS OF THE PROJECT	8
3. MATERIALS AND METHODS	9
3.1. Cell culture	9
3.2. Viability assays.....	9
3.3. Reverse transcriptase-quantitative polymerase chain reaction (RT-qPCR)	9
3.4. Flow cytometry.....	10
3.5. Western blot	10
3.6. Statistical analysis	11
4. RESULTS AND DISCUSSION	12
4.1. RANK and RANKL expression in TNBC cell lines	12
4.2. RANKL/RANK pathway activation by exogenous RANKL.....	14
4.3. RANKL blockade by OPG-Fc.....	15
4.4. Effect of PARPi and PI3Ki in combination with OPG-Fc in TNBC cell viability	16
4.5. Effect of olaparib and alpelisib on downstream molecular targets	18
5. CONCLUSION AND FUTURE PERSPECTIVES	20
REFERENCES.....	21
APPENDIX	34

LIST OF FIGURES

Figure 1.1 Physiological involvement of the RANKL/RANK pathway in mammary epithelial cells....	5
Figure 1.2 Proposed mechanism for progesterone/RANKL-mediated BRCA1 ^{mut} breast cancer carcinogenesis	7
Figure 1.3 The vicious cycle of bone metastases	7
Figure 4.1 Analysis of RANK expression in TNBC cell lines	12
Figure 4.2 Analysis of RANKL expression in TNBC cell lines	13
Figure 4.3 The RANKL/RANK pathway is activated by exogenous RANKL	14
Figure 4.4 OPG-Fc has an inhibitory effect on RANKL binding to RANK	15
Figure 4.5 Cell viability of TNBC cells after single or combinatory treatment with olaparib, alpelisib and OPG-Fc	16
Figure 4.6 Olaparib and alpelisib affect downstream molecular targets	18

LIST OF TABLES

Table S1 BRCA1/2, PTEN and PIK3CA status of the TNBC cell lines in the panel	34
Table S2 mRNA expression data for RANK (TNFRSF11A) and RANKL (TNFSF11) across BC cell lines, retrieved from the Cancer Cell Line Encyclopedia (CCLE).....	34
Table S3 Described IC ₅₀ values for olaparib and alpelisib across the TNBC cell lines.....	35

LIST OF ABBREVIATIONS

ADAM	A disintegrin and metalloproteinase
ADP	Adenosine diphosphate
ALP	Alpelisib
APS	Ammonium persulfate
ATR	Ataxia Telangiectasia and Rad3-related
BC	Breast cancer
BER	Base excision repair
BM	Bone metastasis
BRCA 1/2	Breast cancer 1/2
BSA	Bovine serum albumin
CCLE	Cancer Cell Line Encyclopaedia
cDNA	Complementary deoxyribonucleic acid
Chk1	Checkpoint kinase 1
DC	Dendritic cell
DFS	Disease-free survival
DMBA	Dimethylbenz[a]anthracene
DMEM	Dulbecco's Modified Eagle Medium
DMSO	Dimethyl sulfoxide
DNA	Deoxyribonucleic acid
DSB	Double-strand break
EMA	European Medicines Agency
EMT	Epithelial-mesenchymal transition
ER	Oestrogen receptor
ERK	Extracellular signal-regulated kinase
ETS1	E26 transformation-specific 1
FACS	Fluorescence-activated cell sorting
FBS	Foetal bovine serum
FDA	United States Food and Drug Administration
GAG	Glycosaminoglycans
HER2	Human epidermal growth factor 2
HR	Hormone receptors
HRP	Horseradish peroxidase
HRR	Homologous recombination repair
HRT	Hormone replacement therapy
HT	Hormone therapy
Id2	Inhibitor of DNA binding protein 2
IGF-1	Insulin-like growth factor 1
IKK-α	Inhibitor-kappa B kinase alpha
IL	Interleukin
IκBα	Inhibitor of kappa B alpha
JNK	c-Jun N-terminal kinase
LEC	Luminal epithelial cell
LGR4	Leucine-rich repeat-containing G-protein-coupled receptor 4
LOH	Loss of heterozygosity
MAPK	Mitogen-activated protein kinase
MaSC	Mammary adult stem cell

MEC	Myoepithelial cell
MEM	Minimum Essential Medium Eagle
MITF	Melanocyte inducing transcription factor
MMP	Matrix metalloproteinase
MPA	Medroxyprogesterone acetate
mRNA	Messenger ribonucleic acid
mTOR	Mammalian target of rapamycin
NAD⁺	Nicotinamide adenine dinucleotide
NF-κB	Nuclear factor kappa B
NHEJ	Non-homologous end joining
NTC	No template control
ODF	Osteoclastic differentiation factor
OLA	Olaparib
OPG	Osteoprotegerin
OPGL	Osteoprotegerin ligand
OS	Overall survival
PARP	Poly (adenosine diphosphate-ribose) polymerase
PARPi	Poly (adenosine diphosphate-ribose) polymerase inhibitor
PARylation	Poly adenosine diphosphate ribosylation
PBS	Phosphate buffered saline
pCR	Pathologic complete response
PI3K	Phosphoinositide 3-kinase
PI3Ki	Phosphoinositide 3-kinase inhibitor
PIP2	Phosphatidylinositol (4,5)-bisphosphate
PIP3	Phosphatidylinositol (3,4,5)-triphosphate
PKB	Protein kinase B
PLCγ	Phospholipase gamma
PR	Progesterone receptor
PTEN	Phosphatase and tensin homolog
PTHrP	Parathyroid hormone-related protein
RANK	Receptor activator of nuclear factor kappa B
RANKL	Receptor activator of nuclear factor kappa B ligand
RFS	Relapse-free survival
RNA	Ribonucleic acid
RPMI	Roswell Park Memorial Institute Medium
RT-qPCR	Reverse transcriptase-quantitative polymerase chain reaction
SDS-PAGE	Sodium dodecyl sulphate-polyacrylamide gel electrophoresis
SSB	Single-strand break
TACE	Tumour necrosis factor alpha converting enzyme
TEMED	Tetramethylethylenediamine
TGF-β	Transforming growth factor beta
TNBC	Triple-negative breast cancer
TNF	Tumour necrosis factor
TRAF	TNF-receptor associated factors
TRAIL	Tumour necrosis factor-related apoptosis-inducing ligand
TRANCE	Tumour necrosis factor-related activation-induced cytokine
TRANCE-R	Tumour necrosis factor-related activation-induced cytokine receptor
vWF	von Willebrand factor

1. STATE OF THE ART

Breast cancer (BC) is one of the cancer types with highest worldwide mortality rate, second only to lung cancer. It is the most common form of cancer affecting women¹ and one in every eight women is predicted to develop BC during their lifetime². BC is a multifactorial disease, and its risk factors include genetic predisposition, age over 40 years, early menarche, late menopause, Caucasian race, late childbearing and some hormonal contraception and combined hormone replacement therapy (HRT) regimens³⁻⁵. Less than 25% of BC cases are associated with familial susceptibility, of which around 20% are due to mutations in the BC susceptibility genes *BRCA1* and *BRCA2*⁶. Female *BRCA1* and *BRCA2* mutation carriers have a risk of developing BC by the age of 80 of 72% and 69%, respectively⁷.

BC is genetically and clinically heterogeneous⁸, an aspect which led to its subgrouping to attempt a more specific treatment of patients. The classical immunohistochemical categorisation focuses on the expression of hormone receptors (HR), namely progesterone receptor (PR) and oestrogen receptor (ER), and human epidermal growth factor 2 (HER2, also referred to as ErbB2) amplification⁹. Based on this, BC can be divided into three major subtypes: luminal (ER⁺ and/or PR⁺), which can be further discriminated as luminal A (HER2⁻, Ki-67^{low}) or luminal B (HER2⁻, Ki-67^{high}; or HER2⁺); HER2-overexpressing (ER⁻, PR⁻, HER2⁺); and triple-negative BC (TNBC; ER⁻, PR⁻, HER2⁻)¹⁰⁻¹². This subtyping is of clinical relevance because it determines the possibility to use hormone therapy (HT) or HER2-targeted therapy.

1.1. Triple-negative breast cancer

TNBC is defined by lack of expression of both ER and PR and lack of HER2 amplification. Around 10% to 15% of all BC are classified as triple-negative¹³. TNBC presents an aggressive clinical course and has a worse prognosis with reported higher likelihood of distant recurrence and death within 5 years of diagnosis compared with the other subtypes¹⁴⁻¹⁶. It is associated with higher rates of brain, lung and distant nodal metastases, with bone and liver metastases less common than in non-TNBC¹⁷. The prevalence of TNBC is high amongst younger women, especially of African American descent¹⁸, and around 10% to 15% of TNBC patients carry germline mutations in *BRCA1/2*¹⁹. Furthermore, around 70% of all *BRCA1*-mutated BC are TNBC, contrary to *BRCA2*-mutated BC which are usually ER⁺^{20,21}.

Chemotherapy is the current standard of care for both metastatic and non-metastatic TNBC²²⁻²⁴. TNBC is associated with higher chemosensitivity compared to non-TNBC, which is reflected in the higher pathologic complete response (pCR) rates^{25,26}. However, in patients who did not achieve pCR, it is associated with a poor prognosis because of the high likelihood of early relapse due to acquired resistance to chemotherapeutic drugs^{26,27}. This contrast is termed “triple-negative paradox”.

For this reason, the lack of targeted therapies is a significant constraint to the effective treatment of TNBC patients. An exception are TNBC patients with germline mutations in *BRCA1/2*, tumour suppressor genes which encode proteins with crucial functions in homologous recombination repair (HRR), for whom a targeted approach, poly-(ADP-ribose) polymerase (PARP) inhibition, is available after chemotherapy in the adjuvant and/or metastatic setting²². Moreover, because TNBC is also very heterogeneous, it is more likely that future therapeutic approaches will focus on a specific molecular subtype rather than the whole group of tumours. Several studies have already proposed different molecular classifications for TNBC but they have had little impact in the clinical setting so far²⁸⁻³².

1.1.1. PARP inhibitors (PARPi) in the treatment of TNBC

PARP-1 and PARP-2 are enzymes involved in base excision repair (BER), a DNA damage repair mechanism for DNA single-strand breaks (SSB) predominantly during S-phase, when DNA is exposed for replication³³. In a process termed poly-ADP-ribosylation (PARylation), PARP-1 and PARP-2 act as catalysts of the addition of adenosine diphosphate (ADP) chains to target proteins, with nicotinamide adenine dinucleotide (NAD⁺) as a substrate. PARylation of histones relaxes the chromatin, allowing for the recruitment of DNA repair proteins to sites of damage³⁴. PARPi are small molecules that bind reversibly to the NAD⁺ site of PARP-1 and PARP-2 and prevent PARylation, thus also preventing BER-mediated SSB repair. Moreover, there are evidences that PARPi compromise PARP-1 strand ligation, resulting in continuous turnover of PARP-1 at damage sites and increased sister-chromatid exchange³⁵.

During the S/G2-phase, HRR predominates over non-homologous end joining (NHEJ) regarding DNA double-strand break (DSB) repair because of the presence of a sister chromatid. HRR is regarded as relatively error-proof, contrary to the more error-prone NHEJ. However, some cancers may present impaired HRR, including BC with mutations on *BRCA1* and/or *BRCA2*³⁶. As per the “two-hit” hypothesis formulated by Alfred Knudson³⁷, cancer cells of *BRCA* mutation carriers tend to present loss of the healthy *BRCA1/2* allele through somatic mutation or epigenetic silencing, in a process named loss of heterozygosity (LOH), which represents the second “hit” for tumour suppressor inactivation and impairs HRR of DSB^{38,39}. PARPi are effective in *BRCA1/2*^{-/-} cancer cells via synthetic lethality: inhibition of PARP leads to the accumulation of SSB, whilst preventing their repair and therefore resulting in the formation of DSB; simultaneously, the intrinsic deficiency in HRR presented by cancer cells leads to increased apoptosis due to incorrect DNA repair. A very significant advantage of PARPi-related synthetic lethality resides in its specificity – because non-cancerous cells are heterozygous for *BRCA1/2* mutations, the toxicity is focused on cancer cells⁴⁰.

In the context of BC, the PARPi olaparib (Lynparza[®], AstraZeneca Pharmaceuticals LP) was approved by the US Food & Drug Administration (FDA, in 2018) and the European Medicines Agency (EMA, in 2019) as monotherapy for the treatment of (1) patients with germline *BRCA1/2* mutations presenting HER2⁻ metastatic or locally advanced BC and previously treated with a taxane and an anthracycline (two chemotherapeutic agents) unless not suitable for these treatments; and also (2) patients with HR⁺ BC showing tumour progression on or after endocrine therapy, or otherwise considered unsuitable for said approach. Olaparib had previously been approved, in 2014, for patients with ovarian, fallopian and peritoneal cancer, namely those with *BRCA* mutations, and more recently, in 2020, for patients with metastatic pancreatic cancer presenting germline *BRCA* mutations^{41–43}.

PARP inhibition is the only targeted approach currently approved for use in a subset of TNBC patients with germline *BRCA* mutations. However, other TNBC patients might also benefit from olaparib, namely those presenting somatic *BRCA* mutations⁴⁴ or *BRCA* wild type with gene inactivation through promoter methylation^{45,46}.

1.1.2. PI3K inhibitors (PI3Ki) in the treatment of TNBC

The phosphoinositide 3-kinase (PI3K)/Akt/mammalian target of rapamycin (mTOR) pathway is an important regulator of glucose metabolism and cell growth physiologically triggered upstream through stimulation of receptor tyrosine kinases by growth factors such as insulin⁴⁷. It is the most frequently activated signalling pathway in BC and is mutated in around 40% of TNBC patients⁴⁸.

The *PIK3CA* gene encodes for the p110 α subunit of PI3K α and was reported as the second most common gene mutation amongst TNBC, presenting activation in around 20% to 25% of cases^{49–51}. There are many isoforms of PI3K, which have been divided into three classes (I, II and III) according to structural characteristics and substrate preferences. However, the class I PI3K isoforms, composed of a p85 regulatory subunit and a p110 catalytic subunit^{52,53}, are the most commonly altered in cancer. In cancer, these protein complexes may be mutated so as to activate the signalling pathway, and this can occur either by inactivation of p85 or by overactivation of p110⁵⁴, which is the case of *PIK3CA* mutations.

Phosphatase and tensin homologue (PTEN) is a tumour suppressor gene encoding for a lipid phosphatase that regulates the PI3K/Akt/mTOR pathway through dephosphorylation of phosphoinositide-3,4,5-trisphosphate (PIP3)⁵⁵, opposing PI3K activation of protein serine/threonine kinase Akt (also called protein kinase B, PKB)^{56,57}. Cancer cells benefit from the inactivation of *PTEN*, which can either occur through mutation⁵⁸ or promoter hypermethylation⁵⁹.

Alpelisib (Piqray[®], Novartis Pharmaceuticals Corporation) is a PI3Ki with specificity towards p110 α ⁶⁰, and because it is an isoform-specific inhibitor it presents less off-target adverse effects when compared with pan-PI3K inhibitors⁵⁴. It induces apoptosis through a mechanism involving Akt-dependent induction of Bim, a pro-apoptotic protein, and degradation of Mcl-1, an apoptosis inhibitor⁶¹. Alpelisib was approved by the FDA in 2019 and, very recently, by the EMA (in 2020), in combination with the ER antagonist fulvestrant for the treatment of men and postmenopausal women with HR⁺, HER2⁺ metastatic or locally advanced BC presenting *PIK3CA* mutations, following progression after endocrine monotherapy^{62,63}. Despite clinical data showing no benefit in the addition of a pan-PI3Ki to a chemotherapeutic regimen for TNBC patients^{64,65}, PI3Ki might still have value in combination strategies, since in these studies no selection was made for *PIK3CA*-mutated TNBC.

1.1.3. PARPi and PI3Ki synergy

Interestingly, in 2012 two studies described *in vitro* and *in vivo* synergy between PI3Ki, using buparlisib (NVP-BKM120, a pan-PI3Ki), and PARPi, using olaparib, in *BRCA* competent or mutant TNBC models. PI3K inhibition was accompanied by downregulation of *BRCAl/2* and an increase in DNA damage and PARylation, reflecting an abrogation of HRR and a dependency on PARP activity, thus sensitising cells to PARP inhibition. *BRCA* downregulation appeared to be mediated by an extracellular signal-regulated kinase (ERK)-dependent activation of the transcription factor E26 transformation-specific 1 (ETS1), thereby suppressing gene transcription^{66,67}. Additionally, PI3Ki impair the production of nucleotides required for DNA repair and synthesis, which could also lead to an increased DNA damage when combined with PARPi⁶⁸. Based on these findings, the phase I clinical trial NCT01623349 focused on the combination of olaparib with PI3Ki (buparlisib or alpelisib) for the treatment of patients with recurrent TNBC or high grade serous ovarian cancer. Results from this ongoing study have provided clinical evidence of synergy between olaparib and alpelisib in HRR-proficient ovarian cancer⁶⁹. However, *BRCA*-mutated ovarian cancer showed little improvement from the combination regimen versus olaparib monotherapy. This suggests that the combination regimen could be effective for *BRCA*-competent TNBC, especially in *PIK3CA*-mutated cases.

1.2. RANKL/RANK pathway

Receptor activator of nuclear factor kappa-B ligand (RANKL; also known as osteoprotegerin ligand (OPGL), TNF-related activation-induced cytokine (TRANCE) and osteoclastic differentiation factor

(ODF)⁷⁰⁻⁷³) is a member of the tumour necrosis factor (TNF) superfamily encoded by the *TNFSF11* gene (13q14) that binds to the signalling receptor RANK (*TNFRSF11A* gene, 18q22.1)^{70,74}. RANKL is a homotrimeric type II membrane protein which can occur in three isoforms: RANKL1, the full-length structure; RANKL2, a shorter form missing part of the intra-cytoplasmic domain; and RANKL3, a soluble isoform missing the N-terminal⁷⁵. Besides alternative splicing, membrane-bound RANKL can be shed in a soluble form through proteolysis catalysed by enzymes such as TNF- α -converting enzyme (TACE), A disintegrin and metalloproteinase (ADAM)-10, matrix metalloproteinase (MMP)-7 and MMP-14⁷⁶⁻⁷⁸. RANKL is highly expressed in bone tissue, lymphoid organs and the vascular system, but is also present in other tissues such as the brain, breast, intestine, kidney, liver, lung and skeletal muscle^{79,80}. Its expression can be induced by factors including progesterone, prolactin, parathyroid hormone-related protein (PTHrP), vitamin D3, TNF- α and the interleukins (IL)-11 and 17^{81,82}.

RANK (or TRANCE receptor (TRANCE-R)), on the other hand, is mainly expressed in the bone, although *RANK* messenger RNAs (mRNAs) have been found in other tissues such as breast, liver, prostate and thymus^{74,83}. It is a homotrimeric type I transmembrane protein and, similarly to other members of the TNF receptor superfamily, RANK lacks tyrosine kinase activity, requiring adapter proteins, the TNF receptor-associated factors (TRAF), to be functional⁸⁴. RANK possesses intracellular TRAF binding sites which are able to interact with TRAF-1, 2, 3, 5 and 6⁸⁴⁻⁸⁶. Amongst these, TRAF6 has been shown to have a predominant role in RANK-associated signalling⁸⁷, leading to the activation of several important transduction pathways, namely PI3K/Akt/mTOR; mitogen-activated protein kinases (MAPKs), including p38, c-Jun N-terminal kinase (JNK) and ERK1/2; and Src/phospholipase gamma (PLC γ). This leads to the nuclear translocation of transcriptional activators, primarily nuclear factor kappa-B (NF- κ B) but also others such as Fos/Jun and melanocyte inducing transcription factor (MITF), and results in the transcription of numerous effector genes associated with cell survival, differentiation and proliferation^{86,88,89}. Several RANK alternative splicing isoforms have been described, of which RANK-c stands out due to its ability to counteract RANK-mediated NF- κ B activation^{90,91}.

The RANKL/RANK pathway is negatively regulated by the soluble decoy receptor osteoprotegerin (OPG), encoded by the *TNFRSF11B* gene⁹². Despite its ubiquitous expression, OPG is detected predominantly in bone, immune cells and the vascular system^{83,93}. OPG is able to bind RANKL with an affinity approximately 500 times superior to that of RANK, facilitating the internalization of the cytokine and thus reducing its half-life^{94,95}. OPG can be induced by factors such as oestrogen, IL-4 and transforming growth factor beta (TGF- β)⁹⁶, besides being controlled by many ligands including TNF-related apoptosis-inducing ligand (TRAIL)⁹⁷, glycosaminoglycans (GAGs)⁹⁸ and von Willebrand factor (vWF)⁹³, all of which will influence the inhibitory effect. Moreover, in 2016, Luo *et al.* identified a new RANKL receptor, leucine-rich repeat-containing G-protein-coupled receptor 4 (LGR4), involved in a feedback loop that, like OPG, contributes to the control of RANKL activity⁹⁹.

1.2.1. Physiological functions

The RANKL/RANK axis was originally discovered to mediate T-cell and dendritic cell (DC) communication. Activated T-cells express RANKL, which interacts with RANK expressed on DCs, increasing their antigen-presenting capacity. This leads to an increase in the number of antigen-specific T-cells and enhances memory T-cell immunity⁷⁴. Moreover, RANKL acts as a specific survival factor for DCs⁷² and is essential for the development of lymph nodes¹⁰⁰ and thymic function¹⁰¹.

However, its most prominent role is in the bone. It was discovered to be crucial for the regulation of bone formation and resorption, maintaining bone homeostasis¹⁰⁰. Osteoblasts are bone-forming mononucleated cells of mesenchymal origin, which, together with osteocytes, chondrocytes and stromal cells, express RANKL. RANKL binds to its cognate receptor RANK expressed in osteoclast precursor cells, of hematopoietic origin, thereby inducing osteoclast survival and osteoclastogenesis, and subsequent osteoclast-mediated bone resorption. OPG, also produced by osteoblasts, inhibits bone resorption and, therefore, a balanced OPG:RANKL ratio contributes to bone health, avoiding osteoporosis (OPG<RANKL), commonly seen in postmenopausal women, and osteopetrosis (RANKL>OPG)^{70,71,100,102}.

The RANKL/RANK pathway has also been associated with other physiological processes, e.g. in the central nervous system¹⁰³ and in the breast, where RANKL was discovered to be a crucial regulator of the proliferation and differentiation of mammary epithelial cells especially during pregnancy, driving the morphogenesis of the lactating mammary gland⁸¹. The mammary epithelium comprises an outer basal layer containing myoepithelial cells (MECs) and adult mammary stem cells (MaSCs), and an inner luminal layer consisting of luminal epithelial cells (LECs). Whereas LECs line the ducts and alveoli and are involved in milk synthesis and secretion during lactation, MECs facilitate milk excretion due to their contractile phenotype and MaSCs are able to regenerate the entire mammary epithelial tree due to their self-renewable and multipotent potential^{104,105}. Progesterone, a steroid hormone presenting highest levels during the luteal phase of the menstrual cycle and throughout pregnancy¹⁰⁶, is known to have mitogenic effects on the mammary epithelial cells (**Figure 1.1a**) through its binding to the PR in PR⁺ LECs, upregulating RANKL expression due to mRNA stabilisation¹⁰⁷. The newly synthesised RANKL acts in LECs, which are RANK⁺, further upregulating RANKL; and in a paracrine fashion on the HR⁻ MECs and MaSCs, also RANK⁺. Upon RANKL-RANK binding, two main pathways are triggered on mammary epithelial cells (**Figure 1.1b**). On one hand, it leads to the activation of inhibitor kappa-B kinase alpha (IKK- α), which induces proteasome degradation of inhibitor of kappa-B alpha (I κ B α). Its dissociation from NF- κ B allows the latter to migrate to the nucleus and induce transcription of cyclin D1, involved in the G1 phase of the cell cycle. On the other hand, it also promotes the nuclear translocation of inhibitor of DNA binding protein 2 (Id2), a transcriptional regulator which downregulates p21, a cell cycle inhibitor^{108–110}. Additionally, the binding of RANKL to RANK expressed in the surface of basal cells upregulates RANK expression, notably contributing even more to the proliferation and expansion of MaSCs¹¹¹. Besides progesterone, prolactin and PTHrP can also promote RANKL expression in the mammary epithelium⁸¹.




Figure 1.1 Physiological involvement of the RANKL/RANK pathway in mammary epithelial cells. (a) Progesterone binds to PR in PR⁺ luminal epithelial cells and upregulates RANKL expression, which in turn activates RANK in RANK⁺ luminal and basal cells, promoting proliferation and survival. (b) RANKL binding to RANK in mammary epithelial cells triggers two main cascades leading to NF- κ B-dependent cyclin D1 transcription and Id2-mediated p21 downregulation. Adapted from Infante *et al.*, 2019¹¹².

1.2.2. Involvement in cancer

Numerous studies have demonstrated RANK and/or RANKL expression in different tumours^{113–115}, and others have even described this expression as predictive of poor prognosis^{116–118}. Together with observations regarding physiological functions involving cell survival and proliferation, this hints at a potential tumorigenic involvement of the RANKL/RANK/OPG axis. Rightfully so, this pathway has been implicated in every stage of cancer development, promoting events such as epithelial-mesenchymal transition (EMT) and stemness, cell migration, neovascularization of tumours, establishment of distant metastases, and even promoting immunosuppressive responses^{119–125}.

The Women's Health Initiative and the Million Women Study showed an increased BC risk associated with HRT and contraceptives containing synthetic progesterone derivatives (progestins) such as medroxyprogesterone acetate (MPA)^{4,5}. A 2010 study by Schramek *et al.* showed a significant induction of RANKL expression in mammary epithelial cells of female mice treated with MPA and 7,14-dimethylbenz[a]anthracene (DMBA), a carcinogen. This resulted in marked cell proliferation and eventually BC tumorigenesis. However, genetic inactivation and deletion of RANK in the mammary epithelial cells prevented the MPA-associated proliferation and delayed the onset and incidence of MPA-driven BC¹²⁶. These results show an involvement of the RANKL/RANK pathway in progesterone and progestin-driven breast carcinogenesis, an observation further supported by Gonzalez-Suarez *et al.* in the same year¹²⁷. In accordance, BC is one of the cancers in which RANK and RANKL expression has been documented. Moreover, RANK overexpression in normal human mammary epithelial cells was shown to induce mammary gland reconstitution, EMT, migration, stemness and anchorage-independent growth, contributing to cell transformation and metastasis¹²⁵. A posterior study in mice found that the activation of RANK signalling led to expansion of the luminal and basal mammary compartments, resulting in the accumulation of MaSCs, luminal and bipotent progenitors. This impairment of mammary cell fate eventually resulted in hyperplasia and tumorigenesis¹²⁸.

In 2016, two studies identified a crucial role for the RANKL/RANK pathway in the tumorigenesis of *BRCA1* mutation-driven breast cancer, since *RANK* deletion and the inhibition of the pathway appeared to prevent the development of the disease^{129,130}. Notably, since *BRCA1* mutations are commonly associated with TNBC^{20,21}, a correlation between TNBC and the RANKL/RANK pathway was established. The description of an association between RANK/RANKL dual expression in TNBC with a worse prognosis, when compared to patients with RANK⁺, RANKL⁻ TNBC, strengthened this observation¹¹⁶. One of the proposed mechanisms behind BC tumorigenesis involving *BRCA1* mutations and the RANKL/RANK pathway (**Figure 1.2**) is the following: (i) mutation carriers are prone to LOH, spontaneously losing the remaining wild type *BRCA1* allele; (ii) the resulting HRR-incompetent *BRCA1*-deficient cells present increased genomic instability and DNA damage; (iii) during pregnancy, the menstrual cycle or HRT, progesterone or progestins stimulate the proliferation of RANK⁺ luminal progenitor cells via RANKL/RANK; (iv) as a result, the cells acquire genetic mutations, e.g. in the *TP53* gene, culminating in uncontrolled proliferation and development of BC⁸².




Figure 1.2 Proposed mechanism for progesterone/RANKL-mediated *BRCA1*^{mut} breast cancer carcinogenesis. Mammary epithelial cells of *BRCA1* mutation carriers may present impaired homologous recombination due to loss of the healthy *BRCA1* allele. The subsequent genomic instability increases the chances of acquiring mutations, an event which is further promoted by progesterone/progestin stimuli. This way, RANKL/RANK-mediated proliferation of *BRCA1* deficient cells may lead to the generation of driver mutations, originating breast cancer. Adapted from Rao *et al.*, 2018¹³¹.

Besides primary cancers, the RANKL/RANK pathway is very important in the context of bone metastases (BM), with bone being the most common site of relapse in BC patients. Indeed, this pathway has been shown to promote BC-associated BMs¹³². Circulating BC cells, originating from the primary tumour, are attracted to the bone by chemokines¹³³. Several bone cells, such as osteoblasts, osteocytes and stromal and lining cells, secrete RANKL, activating osteoclast-mediated bone resorption. Bone resorption releases growth factors including TGF- β and insulin-like growth factor 1 (IGF-1), which in turn stimulate cancer cell proliferation and production of several cytokines like PTHrP and IL-6. In turn, these cytokines bind to stromal cells, osteoblasts and other RANKL-producing cells to upregulate RANKL expression. This establishes a “vicious cycle” of osteolytic damage (**Figure 1.3**). Even though osteoblastic (associated with enhanced bone deposition) or mixed bone lesions may also occur, the described osteolytic metastases (associated with increased bone degradation) account for the majority of metastatic BC cases^{134–137}. BMs are a good example of the “seed and soil” hypothesis, where Stephen Paget suggested that when tumour cells leave the primary tumour, they usually target organs with favourable characteristics¹³⁸.




Figure 1.3 The vicious cycle of bone metastases. RANKL produced by stromal cells and osteoblasts promotes osteoclast-mediated bone resorption, releasing molecules that promote cancer cell proliferation. In turn, cancer cells secrete factors which enhance RANKL production even further, giving continuity to the cycle. Adapted from de Groot *et al.*, 2018¹³⁷.

1.2.3. Pathway blockade

The involvement of the RANKL/RANK pathway in several aspects of cancer biology suggests a possible therapeutic potential in its inhibition. Between 2010 and 2011, denosumab, a fully humanised IgG2 anti-RANKL monoclonal antibody, was approved by the FDA and the EMA for clinical use. It was proven effective in breast and prostate cancer patients receiving hormone ablation therapy, and in the prevention of skeletal-related events in multiple myeloma patients and patients with BMs from solid tumours. By blocking the binding of RANKL to RANK, denosumab is able to reduce bone resorption, mimicking the action of OPG^{139–142}.

Evidence suggest an anti-tumour effect for RANKL blockade either through direct inhibition of the pathway in RANK and RANKL-expressing tumours such as lung cancer¹⁴³; by osteoclast-dependent modulation of the bone microenvironment as happens in BMs¹⁴³; or even through the effect of the inhibition on non-cancerous cells, such as immune cells¹⁴⁴. It is the case of the preclinical works that described the involvement of the RANKL/RANK pathway in *BRCA1* mutation-driven BC, in which, as previously mentioned, RANKL inhibition was able to reduce the expansion of mammary progenitors and prevent BC development^{129,130}. This hints at a possible therapeutic application of RANKL blockade in TNBC chemoprevention, and possibly also in combination therapies.

2. AIMS OF THE PROJECT

As previously mentioned, since the RANKL/RANK pathway has been implicated in the tumorigenesis of TNBC, pharmacological inhibition of this pathway could be of clinical relevance beyond bone metastatic disease. We hypothesised it could synergise with PARP and/or PI3K inhibitors, which are not currently in use for all TNBC patients.

Therefore, the main objective of this project was to verify whether RANKL blockade could improve the efficacy of PARPi and PI3Ki, in combination therapy, in *in vitro* TNBC models.

Besides sensitisation of cells to other drugs and/or increasing efficacy in already sensitive cells, a combination therapy based on RANKL blockade may include some advantages that have been associated with combination regimens, including minimizing toxicity and allowing for drug repurposing, such as proposed in the current project, to allow for a faster introduction into the clinical setting^{145,146}.

3. MATERIALS AND METHODS

3.1. Cell culture

The human TNBC cell line panel was selected according to the mutational status of *BRCA1/2*, *PIK3CA* and *PTEN* (**Table S1**), which may influence drug response. The MDA-MB-231^{GFP+/Luc+} cell line (hereby referred as MDA-MB-231) was kindly provided by Sérgio Dias (Instituto de Medicina Molecular, Lisbon, Portugal); the BT-20 cell line was purchased from ATCC (HTB-19TM); and the MDA-MB-468 and HCC1937 cell lines were generously supplied by Rita Fior (Champalimaud Centre for the Unknown, Lisbon, Portugal). All cell lines were tested for *Mycoplasma* contamination.

The cell lines were cultured in media supplemented with 10% (v/v) Foetal Bovine Serum (FBS, Gibco) and 1% (v/v) Penicillin/Streptomycin (Pen/Strep: 10.000 U/mL Penicillin, 10.000 µg/mL Streptomycin; Gibco). Dulbecco's Modified Eagle Medium (DMEM) 1X containing D-Glucose, L-Glucose and Pyruvate (Gibco) was used for the MDA-MB-231 and MDA-MB-468 cell lines; Minimum Essential Medium Eagle (MEM) 1X containing Earle's Salts and L-Glutamine (Gibco) for the BT-20 cell line; and Roswell Park Memorial Institute Medium (RPMI) 1640 1X (Gibco) containing L-Glutamine for the HCC1937 cell line. Cells were maintained in T25 or T75 cell culture flasks (Thermo Scientific) at 37°C with 5% CO₂ and humidified atmosphere. Cells were used at low passages and media was changed every 2 to 3 days.

3.2. Viability assays

To assess viability, cells were seeded in 96-well plates (Costar or TPP) with a density of 10⁴ cells/well and treated with different concentrations of olaparib (#10-2154, Focus Biomolecules), alpelisib (#ENZ-CHM214, Enzo Life Sciences) and OPG-Fc (Amgen, Inc.). Olaparib was solubilised in absolute ethanol; alpelisib in DMSO; and OPG-Fc in PBS. After incubation for 72h under standard conditions, cell viability was assessed using the Alamar Blue viability reagent (Invitrogen). 1:10 (v/v) reagent was added to the media of each well and, after a 2h incubation at 37°C, fluorescence (excitation 560 nm / emission 590 nm) was read in a microplate reader (Infinite M200, TECAN). Every experiment was done in triplicate or more, with six replicates per experiment.

3.3. Reverse transcriptase-quantitative polymerase chain reaction (RT-qPCR)

For *RANK* and *RANKL* mRNA expression analysis, total RNA was extracted with the NZY Total RNA Isolation kit (Nzytech), according to manufacturer's instructions, and quantified using a spectrophotometer (NanoDrop 1000, Thermo Fisher). Total RNA was treated with RQ1 RNase-free DNase (Promega) for 30 min at 37°C, in accordance with manufacturer instructions. 480 ng of DNase I-treated RNA were used for complementary DNA (cDNA) synthesis, using the NZY M-MuLV First-Strand cDNA Synthesis kit (Nzytech), Oligo(dT)₁₈ primer mix (Nzytech) and a thermal cycler (C1000, Bio-Rad). The resulting cDNAs were amplified by real-time PCR with TaqMan Gene Expression Master Mix (Nzytech) and specific primers including: *TNFRSF11A* (#Hs00921372_m1, Applied Biosciences), *TNFSF11* (#Hs00243522_m1, Applied Biosciences) and *GAPDH* (#PPH00150F, SA Biosciences). No template control (NTC) and RNA template controls were included. Every cDNA was tested in triplicate in 384-well PCR plates (Applied Biosystems) and amplification was performed in a Real-Time PCR System (ViiA 7, Applied Biosystems). Gene expression was averaged and normalised using the selected housekeeping gene, glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*), and relative mRNA expression is presented as 2^{-ΔCt}.

3.4. Flow cytometry

To evaluate RANK expression by flow cytometry, cells were resuspended and centrifuged twice for 5 min at 200 x g and 4°C, with a cold PBS wash in between. The pellet was resuspended in fluorescence-activated cell sorting (FACS) buffer (2% FBS, 1X PBS) and divided into three tubes: (A) one for the desired staining; (B) one for the secondary antibody only; and (C) one for the unstained control. In tube A, cells were incubated in the dark for 30 min at 4°C with 5 µL mouse monoclonal anti-RANK antibody (#M331, Amgen Inc.). After washing again with cold PBS, tubes A and B were incubated, also in the dark, for 30 min with Cy5-conjugated AffiniPure goat anti-mouse IgG (#115-175-205, Dianova) diluted 1:100 (v/v) in FACS buffer. Tubes A and B were then washed in FACS buffer with a similar centrifugation as before and all pellets, including from tube C, were resuspended in FACS buffer. Data acquisition was performed in a flow cytometer (LSRFortessa, BD Biosciences) and further analysis was made with the FlowJo vX.0.7 software.

3.5. Western blot

To analyse RANKL/RANK pathway activation with or without OPG-Fc blockade, 2×10^5 cells/well were seeded in 6-well plates (Costar) for 24h and serum-starved in 0.1% FBS, 1% Pen/Strep media for another 24h. The cells were then stimulated for 20 and 60 min with media containing 1 µg/mL human RANKL (hRANKL) (Amgen Inc.) and 60 min with neutralisation media. Two neutralisation media were prepared and incubated for 1h at 37°C: 1 µg/mL hRANKL with 2.5 µg/mL OPG-Fc; and 1 µg/mL hRANKL with 2.5 µg/mL mouse monoclonal anti-hRANKL (#MAB626, R&D Systems; hereby referred as MAB626).

To assess the molecular effect of PARP inhibition using olaparib and PI3K inhibition using alpelisib, 4×10^5 cells/well were plated in 6-well plates and treated for 72h with 10 µM olaparib, 20 µM alpelisib, 100 ng/mL OPG-Fc and all possible combinations.

Proteins were extracted with RIPA buffer (Sigma) with 1:100 (v/v) protease inhibitor cocktail (Cell Signaling) and 1:100 (v/v) phosphatase inhibitor cocktail (Sigma) (herein referred as RIPA buffer), according to manufacturer's instructions. Protein was quantified with the Pierce BCA Protein Assay kit (Thermo Scientific), in accordance with manufacturer's instructions. Protein concentration was calculated using an albumin standard curve and protein extracts were stored at -20°C until usage.

2X or 4X sample buffer with 5% β-mercaptoethanol was added to the extracts before a 10-minute incubation at 95°C. Proteins were resolved on 10% or 12% SDS-PAGE gels and the NZYColour Protein Marker II (Nzytech) was used. Proteins were transferred to nitrocellulose membranes (Invitrogen) using a dry blotting system (iBlot 2, Invitrogen) and visualised with Ponceau S (Sigma) staining after transfer. Membranes were blocked for 1h in PBST (0.05% Tween 20, 1X PBS) 5% (m/v) Albumin Bovine Fraction V (BSA) (Nzytech), hereby designated as BSA; or in PBST 5% (m/v) non-fat dry milk (Molico), hereby designated as milk. Incubation with primary antibodies was overnight at 4°C and with secondary antibodies was 2h at room temperature. The following primary antibodies were used: mouse monoclonal anti β-Actin antibody (1:25000 in milk; #ab6276; Abcam), rabbit monoclonal anti-NF-kB p65 (D14E12) (1:1000 in BSA; #8242, Cell Signaling), rabbit monoclonal anti-Phospho-NF-kB p65 (Ser536) (93H1) (1:1000 in BSA; #3033, Cell Signaling), mouse monoclonal anti-IκBα (L35A5) (1:1000 in BSA; #4814, Cell Signaling), rabbit monoclonal p44/42 MAPK (Erk1/2) (137F5) (1:1000 in BSA; #4695, Cell Signaling), rabbit polyclonal anti-Phospho-ERK1/2 (Thr 202/Tyr 204) (1:1000 in BSA; #sc-16982, Santa Cruz Biotechnology), rabbit monoclonal anti-Akt (pan) (11E7) (1:1000 in BSA; #4685, Cell Signaling), rabbit monoclonal anti-Phospho-Akt (Ser473) (D9E) (1:1000 in BSA; #4060,

Cell Signaling), rabbit polyclonal anti-hsRANKL (1:5000 in BSA; #500-P133, PeproTech), rabbit monoclonal anti-Phospho-Chk1 (Ser345) (133D3) (1:500 in BSA; #2348, Cell Signaling), mouse monoclonal anti-S6 (H-4) (1:500 in BSA; sc-74576, Santa Cruz Biotechnology), rabbit polyclonal anti-Phospho-S6 (Ser240/244) (1:1000 in BSA; #2215, Cell Signaling). Also, the following horseradish peroxidase (HRP)-conjugated specific secondary antibodies were used: anti-mouse-HRP IgG (1:5000 in milk; #7076, Cell Signaling), and anti-rabbit-HRP IgG (1:5000 in milk; #7074, Cell Signaling).

Target proteins were detected using the Novex ECL HRP Chemiluminescent Substrate Reagent kit (Invitrogen) in a blot and gel imager (Amersham Imager 680, GE Healthcare). Some membranes were stripped with membrane stripping solution (Nzytech) and reused. Band intensity was calculated using the ImageJ 1.48v software and normalised for β -actin and, when applicable, to total protein.

3.6. Statistical analysis

GraphPad Prism software version 8.4.3 was used for data analysis and construction of graphics. For the cell viability assays, one-way ANOVA was performed. Values are presented as mean \pm SEM and $p < 0.05$ was considered for statistical significance.

4. RESULTS AND DISCUSSION

4.1. RANK and RANKL expression in TNBC cell lines

TNBC has been shown to express RANK and RANKL¹¹⁶. We started by quantifying endogenous RANK expression across a TNBC cell line panel, by RT-qPCR and flow cytometry (**Figure 4.1**); and endogenous RANKL expression by RT-qPCR and western blot (**Figure 4.2**).

In accordance with data available in the Broad Institute Cancer Cell Line Encyclopedia (CCLE) (**Table S2**), MDA-MB-468 and MDA-MB-231 cell lines presented the highest *RANK* mRNA expression amongst the panel, with HCC1937 and BT-20 cell lines presenting lower expression (**Figure 4.1A**). RANK protein expression was further confirmed through surface protein staining (**Figure 4.1B**).




Figure 4.1 Analysis of *RANK* expression in TNBC cell lines. (A) RT-qPCR of *RANK* in HCC1937, BT-20, MDA-MB-231 and MDA-MB-468 cells. Relative mRNA expression was calculated as $2^{-\Delta C_t}$ using the housekeeping gene *GAPDH* for normalisation. (B) Flow cytometry of RANK in the TNBC panel.

Regarding RANKL expression, HCC1937 cells presented the highest *RANKL* mRNA levels (**Figure 4.2A**), again in agreement with data from the CCLE (**Table S2**). However, we did not observe significant differences in protein expression assessed by western blot, even in comparison to MCF-7 luminal BC cells, known to express RANKL (**Figure 4.2B**). RANKL is a homotrimeric protein which can also be shed in a soluble form. The western blot for RANKL presents multiple bands that correspond to multimeric forms of higher molecular weight, which could include dimers and trimers, whereas bands around 35 kDa are possibly monomeric forms of RANKL. Therefore, bands at lower molecular weights should correspond to its soluble form, considering soluble RANKL possesses a shorter structure⁷⁵.




Figure 4.2 Analysis of RANKL expression in TNBC cell lines. (A) RT-qPCR of *RANKL* in HCC1937, BT-20, MDA-MB-231 and MDA-MB-468 cell lines. Relative mRNA expression was calculated as $2^{-\Delta C_t}$ using the housekeeping gene *GAPDH* for normalisation. (B) Western blot of RANKL in the selected TNBC panel with β -actin as loading control; 100 ng of hRANKL were used as a positive control; and protein extract from MCF-7 cells was used as a positive luminal BC control.

TNBC have been reported to express higher levels of surface RANK and *RANK* mRNA compared to non-TNBC, and dual expression of RANK and RANKL was associated with worse overall survival (OS) and relapse-free survival (RFS; also called disease-free survival, DFS) compared to RANK single expression^{16,116}. Cell lines with highest expression of these two molecules might be more susceptible to inactivation of the RANKL/RANK pathway, namely through RANKL blockade.

4.2. RANKL/RANK pathway activation by exogenous RANKL

To assess the possibility of an inhibitory pharmacological approach via RANKL blockade, activation of the RANKL/RANK pathway by exogenous RANKL was confirmed by western blot in serum-starved cell lines of the selected TNBC panel (**Figure 4.3**).




Figure 4.3 The RANKL/RANK pathway is activated by exogenous RANKL. Phosphorylation of downstream targets of the RANKL/RANK pathway and degradation of IκBα were assessed by western blot upon stimulation with 1 μg/mL hRANKL for 20 or 60 minutes. β-actin was included as loading control. Band intensity was quantified with the ImageJ software and ratios to loading control or total protein were calculated.

Following RANK activation by RANKL, IKK-α is phosphorylated and induces the degradation of IκBα by the proteasome, ultimately allowing for NF-κB translocation into the nucleus, a process requiring NF-κB subunit p65 phosphorylation¹⁰⁸. Besides signalling through the NF-κB pathway, RANK also activates the PI3K/Akt/mTOR pathway via TRAF6 and c-Src, leading to the phosphorylation and activation of Akt⁸⁹. Additionally, the MAPK/Erk pathway is also activated downstream of RANK¹⁴⁷. Accordingly, we evaluated the activation of RANK pathway by exogenous RANKL by assessing the levels of total IκBα, phospho (p)-NF-κB p65 subunit, phospho-Akt and phospho-Erk. Following RANKL stimulus, a reduction in IκBα and an increase in p-p65, p-Akt and p-Erk were expected.

The results reflect the dynamics of the pathway activation and downstream molecular cascade in the different cell lines, with different activation rates and sustained activation times before returning to basal levels. This justifies why not every variation was observed in every condition of the two stimulation timepoints. Since we confirmed pathway activation by exogenous RANKL, next we aimed to test the effect of an inhibitory approach via RANKL blockade.

4.3. RANKL blockade by OPG-Fc

RANKL inhibition can be achieved using OPG-Fc, a recombinant protein resulting of truncated OPG fused to the Fc domain of a human IgG1¹⁴⁸. OPG-Fc lacks the heparin-binding and death homology domains of OPG but has an increased circulating half-life. OPG-Fc is used in animal models of RANKL blockade since denosumab, the monoclonal antibody against RANKL, does not recognize murine RANKL.

To confirm that OPG-Fc effectively inhibits RANKL-mediated RANK activation, serum-starved MDA-MB-231 cells were incubated with hRANKL, previously incubated with OPG-Fc or not, and pathway activation was assessed by quantification of p-Erk and p-Akt by western blot (**Figure 4.4**). MAB626, a monoclonal antibody that effectively blocks RANKL¹⁴⁹, was used as control.




Figure 4.4 OPG-Fc has an inhibitory effect on RANKL binding to RANK. MDA-MB-231 cells were treated for 20 minutes with 1 µg/mL RANKL ± 2.5 µg/mL OPG-Fc or MAB626. Downstream targets of the RANKL/RANK pathway were analysed by western blot with β-actin as loading control.

A decrease in Akt and Erk phosphorylation was observed in cells exposed to neutralised RANKL, in comparison to RANKL stimulation, confirming that OPG-Fc impaired RANKL binding to RANK to some extent.

4.4. Effect of PARPi and PI3Ki in combination with OPG-Fc in TNBC cell viability

Since the RANKL/RANK pathway was shown to be activated by exogenous RANKL in the selected cell lines, and OPG-Fc was able to abrogate this activation, next we aimed to test the combinatory effect of RANKL blockade plus PARP and/or PI3K inhibition.

The effect of the PARPi olaparib, the PI3Ki alpelisib and OPG-Fc was assessed by measuring cell viability 72 hours after incubation with the indicated drugs (**Figure 4.5A**). Overall, the results were in line with described IC₅₀ values for olaparib and alpelisib (**Table S3**), with exception of BT-20 cells treated with olaparib, which presented a higher resistance than reported. Regarding OPG-Fc, cell viability showed little variation with increasing dose, suggesting OPG-Fc alone is not able to affect cell viability.




Figure 4.5 Cell viability of TNBC cells after single or combinatory treatment with olaparib, alpelisib and OPG-Fc. (A) 72-hour cell viability assays were conducted in HCC1937, BT-20, MDA-MB-231 and MDA-MB-468 cell lines upon treatment with different concentrations of olaparib (OLA), alpelisib (ALP) and OPG-Fc. (B) Cell viability was evaluated in cells treated with the indicated combinations. Results are the mean of at least three independent experiments and are presented as mean \pm SEM. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$; ns, not significant. One-way ANOVA test was performed using the GraphPad software.

Interestingly, the most sensitive cell line to olaparib was the BRCA-competent MDA-MB-468 cell line. Multiple factors beyond *BRCA* mutations may confer sensitivity to PARP inhibition, namely those which lead to defects in HRR¹⁵⁰. For example, cell lines with wild type *BRCA* may undergo *BRCA* allelic loss, thus becoming more sensitive to PARPi¹⁵¹. Inversely, HCC1937 cells, which harbour a mutation in *BRCA1* (**Table S1**), were quite resistant to olaparib. Resistance to PARPi is a major problem in the clinical setting and can be due mainly to four mechanisms: increased drug efflux, decreased PARP trapping, restoration of HRR and restoration of stalled replication fork stability¹⁵².

Regarding PI3K inhibition with alpelisib, the BT-20 cell line, which has two heterozygous mutations in *PIK3CA* (**Table S1**), presented the highest sensitivity to alpelisib, followed by MDA-MB-231 cells. On the other hand, MDA-MB-468 and HCC1937 were more resistant to alpelisib, in concordance with the fact that *PTEN* mutations, that both present, may confer resistance to PI3K inhibition¹⁵³. In addition, BC cells may adopt resistance mechanisms such as increased dependency on other PI3K isoforms, e.g. cells may rapidly restore PI3K signalling through activation of p110 β ¹⁵⁴.

As previously mentioned, a synergy between PI3Ki and PARPi was described *in vitro* for BC cells^{66,67}. Taking this discovery into consideration, the combinatory approaches pertain three main comparisons: olaparib versus olaparib and OPG-Fc; alpelisib versus alpelisib and OPG-Fc; and olaparib and alpelisib versus olaparib, alpelisib and OPG-Fc. However, in neither of these conditions was the addition of OPG-Fc associated with a statistically significant decrease in cell viability (**Figure 4.5B**). This indicates that there is no synergic effect between RANKL blockade with OPG-Fc and either olaparib and/or alpelisib *in vitro*.

Moreover, combination of olaparib and alpelisib in the indicated dosages did not present a statistically significant difference in cell viability compared to the drug with lowest associated viability – which was alpelisib for all cell lines. Despite this observation, HCC1937 and MDA-MB-468 cells presented a slight reduction in viability after treatment with the olaparib/alpelisib double combination.

4.5. Effect of olaparib and alpelisib on downstream molecular targets

Finally, to confirm that the reduction in cell viability was dependent of effective targeting by the drugs, namely PARP by olaparib or PI3K by alpelisib, HCC1937 and MDA-MB-468 cells were treated as before, and the specific targets analysed by western blot (**Figure 4.6**).




Figure 4.6 Olaparib and alpelisib affect downstream molecular targets. (A) HCC1937 and (B) MDA-MB-468 cell lines were treated with 100 ng/mL OPG-Fc, 10 μ M olaparib, 20 μ M alpelisib or derived combinations for 72 hours. Using β -actin as loading control, olaparib or alpelisib-related targets were evaluated via western blot. Band intensity was quantified with the ImageJ software and ratios to loading control or total protein were calculated.

Because alpelisib is a PI3Ki, it contributes to the inhibition of the PI3K/Akt/mTOR pathway, and therefore a reduction in phospho-Akt was expected following treatment. Curiously, whilst no noticeable change was seen on MDA-MB-468 cells, the opposite was observed in HCC1937 cells, i.e. presence of alpelisib promoted Akt phosphorylation. An *in vitro* 2018 work by Clement *et al.* uncovered a resistance mechanism to PI3Ki consisting of E3 ubiquitin ligase Skp2-dependent reactivation of Akt. According to this research, inhibition or depletion of PI3K could lead to increased Akt phosphorylation – an event restricted to a subset of TNBC cell lines, in which HCC1937 are included¹⁵⁵. The observed increase in

p-Akt in cells treated with alpelisib is likely due to this mechanism, which means that, albeit unexpectedly, a molecular impact of alpelisib is visible. The S6 ribosomal protein is also a downstream target of the PI3K pathway, with pathway activation promoting its phosphorylation. Thus, it was also expected that upon PI3K inhibition phospho-S6 levels would diminish, although no variation in p-S6 could be observed regarding alpelisib treatment.

On the other hand, cells treated with olaparib are expected to present higher levels of phospho-checkpoint kinase 1 (Chk1). The kinase Chk1 is activated via phosphorylation downstream of Ataxia Telangiectasia and Rad3-related (ATR) following DNA damage or replication stress, arresting cell cycle progression to allow for DNA repair. Indeed, increase in p-Chk1 was observed for both HCC1937 and MDA-MB-468 cells upon olaparib treatment.

These results indicate that both olaparib and alpelisib are affecting molecular targets in TNBC cells, although to different extents, an event concomitant with their effect on cell viability. This allows us to exclude drug inefficacy as a possible explanation for the lack of synergy in the drug combinations. There are, however, limitations to this study, as analysis of downstream molecular targets was performed in a single experiment.

5. CONCLUSION AND FUTURE PERSPECTIVES

TNBC is an aggressive subtype of BC associated with poor clinical outcome. One significant constraint to the treatment of patients is the lack of targeted therapies, which would allow a more specific, and likely effective, approach. The discovery of a link between the RANKL/RANK pathway and TNBC tumorigenesis allowed for the investigation of innovative approaches to treat the disease, namely combination therapies including RANK signalling modulation.

In the present project, we hypothesised that RANKL blockade could synergise with the PI3K inhibitor alpelisib and the PARP inhibitor olaparib, broadening their applicability to include some or more TNBC patients, respectively. However, *in vitro* analyses using TNBC cell lines showed no statistically significant effect of combination regimens.

Recently, RANKL blockade was associated with higher pre-clinical efficacy of immune checkpoint inhibitors, namely in combination with PD1/PD-L1 or CTLA4 blockade¹⁵⁶. As of 2020, there are two ongoing clinical trials, KEYPAD (NCT03280667) and CHARLI (NCT03161756), studying a combination of denosumab with immune checkpoint inhibitors in clear cell renal carcinoma and unresectable or metastatic melanoma, respectively. If the pre-clinical evidence is verified in these trials, the same approach should be tested on BC, since the RANKL/RANK pathway has a significant involvement in mammary tissue, both in physiological and oncological contexts. TNBC would be especially attractive in this case since it is more likely to benefit from immune checkpoint blockade than other subtypes due to its higher immunogenicity, higher PD-L1 expression and enrichment in tumour-infiltrating lymphocytes^{157–159}.

Since TNBC patients have very limited therapeutic options, research and development of novel molecules or even drug repurposing are of critical importance. More so considering the high heterogeneity of TNBC, which will likely require subtyping and development of personalised approaches to each subtype. Combination therapies will probably be essential for this process, and although the drug combinations proposed in this project did not show any benefit *in vitro*, RANKL blockade should be studied in depth to direct all of its potential towards the discovery of treatment alternatives for TNBC patients.

REFERENCES

1. Global Cancer Observatory. Published 2018. <https://gco.iarc.fr/>
2. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin.* 2019;69(1):7-34. doi:10.3322/caac.21551
3. Kamińska M, Ciszewski T, Łopacka-Szatan K, Miotła P, Starosławska E. Breast cancer risk factors. *Menopausal Rev.* 2015;3:196-202. doi:10.5114/pm.2015.54346
4. Million Women Study Collaborators. Breast cancer and hormone-replacement therapy in the Million Women Study. *The Lancet.* 2003;362(9382):419-427. doi:10.1016/S0140-6736(03)14065-2
5. Writing Group for the Women's Health Initiative Investigators. Risks and Benefits of Estrogen Plus Progestin in Healthy Postmenopausal Women: Principal Results From the Women's Health Initiative Randomized Controlled Trial. *JAMA J Am Med Assoc.* 2002;288(3):321-333. doi:10.1001/jama.288.3.321
6. Balmaña J, Díez O, Rubio IT, Cardoso F. BRCA in breast cancer: ESMO Clinical Practice Guidelines. *Ann Oncol.* 2011;22:vi31-vi34. doi:10.1093/annonc/mdr373
7. Kuchenbaecker KB, Hopper JL, Barnes DR, et al. Risks of Breast, Ovarian, and Contralateral Breast Cancer for BRCA1 and BRCA2 Mutation Carriers. *JAMA.* 2017;317(23):2402-2402. doi:10.1001/jama.2017.7112
8. Stingl J, Caldas C. Molecular heterogeneity of breast carcinomas and the cancer stem cell hypothesis. *Nat Rev Cancer.* 2007;7(10):791-799. doi:10.1038/nrc2212
9. Williams SL, Birdsong GG, Cohen C, Siddiqui MT. Immunohistochemical detection of estrogen and progesterone receptor and HER2 expression in breast carcinomas: comparison of cell block and tissue block preparations. *Int J Clin Exp Pathol.* 2009;2(5):476-480.
10. Sørlie T, Perou CM, Tibshirani R, et al. Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci U S A.* 2001;98(19):10869-10874. doi:10.1073/pnas.191367098
11. Perou CM, Sørlie T, Eisen MB, et al. Molecular portraits of human breast tumours. *Nature.* 2000;406(6797):747-752. doi:10.1038/35021093
12. Pathmanathan N, Balleine RL. Ki67 and proliferation in breast cancer. *J Clin Pathol.* 2013;66(6):512-516. doi:10.1136/jclinpath-2012-201085
13. Howlader N, Altekruse SF, Li CI, et al. US incidence of breast cancer subtypes defined by joint hormone receptor and HER2 status. *J Natl Cancer Inst.* 2014;106(5). doi:10.1093/jnci/dju055
14. Bardou V-J, Arpino G, Elledge RM, Osborne CK, Clark GM. Progesterone receptor status significantly improves outcome prediction over estrogen receptor status alone for adjuvant endocrine therapy in two large breast cancer databases. *J Clin Oncol Off J Am Soc Clin Oncol.* 2003;21(10):1973-1979. doi:10.1200/JCO.2003.09.099

15. Dent R, Trudeau M, Pritchard KI, et al. Triple-Negative Breast Cancer: Clinical Features and Patterns of Recurrence. *Clin Cancer Res.* 2007;13(15):4429-4434. doi:10.1158/1078-0432.CCR-06-3045
16. Agarwal G, Nanda G, Lal P, et al. Outcomes of Triple-Negative Breast Cancers (TNBC) Compared with Non-TNBC: Does the Survival Vary for All Stages? *World J Surg.* 2016;40(6):1362-1372. doi:10.1007/s00268-016-3422-4
17. Kennecke H, Yerushalmi R, Woods R, et al. Metastatic behavior of breast cancer subtypes. *J Clin Oncol Off J Am Soc Clin Oncol.* 2010;28(20):3271-3277. doi:10.1200/JCO.2009.25.9820
18. Siddharth S, Sharma D. Racial Disparity and Triple-Negative Breast Cancer in African-American Women: A Multifaceted Affair between Obesity, Biology, and Socioeconomic Determinants. *Cancers.* 2018;10(12):514. doi:10.3390/cancers10120514
19. Armstrong N, Ryder S, Forbes C, Ross J, Quek RG. A systematic review of the international prevalence of BRCA mutation in breast cancer. *Clin Epidemiol.* 2019;11:543-561. doi:10.2147/CLEP.S206949
20. Chen H, Wu J, Zhang Z, et al. Association Between BRCA Status and Triple-Negative Breast Cancer: A Meta-Analysis. *Front Pharmacol.* 2018;9:909-909. doi:10.3389/fphar.2018.00909
21. Mavaddat N, Barrowdale D, Andrulis IL, et al. Pathology of breast and ovarian cancers among BRCA1 and BRCA2 mutation carriers: results from the Consortium of Investigators of Modifiers of BRCA1/2 (CIMBA). *Cancer Epidemiol Biomark Prev Publ Am Assoc Cancer Res Cosponsored Am Soc Prev Oncol.* 2012;21(1):134-147. doi:10.1158/1055-9965.EPI-11-0775
22. Cardoso F, Senkus E, Costa A, et al. 4th ESO–ESMO International Consensus Guidelines for Advanced Breast Cancer (ABC 4). *Ann Oncol.* 2018;29(8):1634-1657. doi:10.1093/annonc/mdy192
23. Cardoso F, Kyriakides S, Ohno S, et al. Early breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up†. *Ann Oncol Off J Eur Soc Med Oncol.* 2019;30(8):1194-1220. doi:10.1093/annonc/mdz173
24. Waks AG, Winer EP. Breast Cancer Treatment. *JAMA.* 2019;321(3):288-288. doi:10.1001/jama.2018.19323
25. Liedtke C, Mazouni C, Hess KR, et al. Response to neoadjuvant therapy and long-term survival in patients with triple-negative breast cancer. *J Clin Oncol Off J Am Soc Clin Oncol.* 2008;26(8):1275-1281. doi:10.1200/JCO.2007.14.4147
26. Carey LA, Dees EC, Sawyer L, et al. The Triple Negative Paradox: Primary Tumor Chemosensitivity of Breast Cancer Subtypes. *Clin Cancer Res.* 2007;13(8):2329-2334. doi:10.1158/1078-0432.CCR-06-1109
27. Nedeljković M, Damjanović A. Mechanisms of Chemotherapy Resistance in Triple-Negative Breast Cancer—How We Can Rise to the Challenge. *Cells.* 2019;8(9):957-957. doi:10.3390/cells8090957

28. Prado-Vázquez G, Gámez-Pozo A, Trilla-Fuertes L, et al. A novel approach to triple-negative breast cancer molecular classification reveals a luminal immune-positive subgroup with good prognoses. *Sci Rep*. 2019;9(1):1538-1538. doi:10.1038/s41598-018-38364-y
29. Lehmann BD, Bauer JA, Chen X, et al. Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. *J Clin Invest*. 2011;121(7):2750-2767. doi:10.1172/JCI45014
30. Lehmann BD, Jovanović B, Chen X, et al. Refinement of Triple-Negative Breast Cancer Molecular Subtypes: Implications for Neoadjuvant Chemotherapy Selection. *PloS One*. 2016;11(6). doi:10.1371/journal.pone.0157368
31. Burstein MD, Tsimelzon A, Poage GM, et al. Comprehensive genomic analysis identifies novel subtypes and targets of triple-negative breast cancer. *Clin Cancer Res Off J Am Assoc Cancer Res*. 2015;21(7):1688-1698. doi:10.1158/1078-0432.CCR-14-0432
32. Jézéquel P, Sharif Z, Lasla H, et al. Gene-expression signature functional annotation of breast cancer tumours in function of age. *BMC Med Genomics*. 2015;8(1):80. doi:10.1186/s12920-015-0153-6
33. Heale JT, Ball AR, Schmiesing JA, et al. Condensin I Interacts with the PARP-1-XRCC1 Complex and Functions in DNA Single-Strand Break Repair. *Mol Cell*. 2006;21(6):837-848. doi:10.1016/j.molcel.2006.01.036
34. Schreiber V, Dantzer F, Ame J-C, de Murcia G. Poly(ADP-ribose): novel functions for an old molecule. *Nat Rev Mol Cell Biol*. 2006;7(7):517-528. doi:10.1038/nrm1963
35. Shao Z, Lee BJ, Rouleau-Turcotte É, et al. Clinical PARP inhibitors do not abrogate PARP1 exchange at DNA damage sites in vivo. *Nucleic Acids Res*. Published online September 5, 2020:gkaa718. doi:10.1093/nar/gkaa718
36. Moynahan ME, Jasin M. Mitotic homologous recombination maintains genomic stability and suppresses tumorigenesis. *Nat Rev Mol Cell Biol*. 2010;11(3):196-207. doi:10.1038/nrm2851
37. Knudson AG. Mutation and Cancer: Statistical Study of Retinoblastoma. *Proc Natl Acad Sci*. 1971;68(4):820-823. doi:10.1073/pnas.68.4.820
38. Smith SA, Easton DF, Evans DGR, Ponder BAJ. Allele losses in the region 17q12–21 in familial breast and ovarian cancer involve the wild-type chromosome. *Nat Genet*. 1992;2(2):128-131. doi:10.1038/ng1092-128
39. Gudmundsson J, Johannesdottir G, Bergthorsson JT, et al. Different tumor types from BRCA2 carriers show wild-type chromosome deletions on 13q12-q13. *Cancer Res*. 1995;55(21):4830-4832.
40. Caulfield SE, Davis CC, Byers KF. Olaparib: A Novel Therapy for Metastatic Breast Cancer in Patients With a BRCA1/2 Mutation. *J Adv Pract Oncol*. 2019;10(2):167-174. <http://www.ncbi.nlm.nih.gov/pubmed/31538027>

41. U.S. Food & Drug Administration. FDA approves olaparib for germline BRCA-mutated metastatic breast cancer. Published 2018. <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-olaparib-germline-brca-mutated-metastatic-breast-cancer>
42. European Medicines Agency. Lynparza. Published 2019. <https://www.ema.europa.eu/en/medicines/human/EPAR/lynparza>
43. U.S. Food & Drug Administration. FDA approves olaparib for gBRCAm metastatic pancreatic adenocarcinoma. Published 2020. <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-olaparib-gbrcam-metastatic-pancreatic-adenocarcinoma>
44. Mohyuddin GR, Aziz M, Britt A, et al. Similar response rates and survival with PARP inhibitors for patients with solid tumors harboring somatic versus Germline BRCA mutations: a Meta-analysis and systematic review. *BMC Cancer*. 2020;20(1):507. doi:10.1186/s12885-020-06948-5
45. Esteller M, Silva JM, Dominguez G, et al. Promoter Hypermethylation and BRCA1 Inactivation in Sporadic Breast and Ovarian Tumors. *J Natl Cancer Inst*. 2000;92(7):564-569. doi:10.1093/jnci/92.7.564
46. De La Haba J, Guerrero-Zotano A, Perez-Fidalgo JA, et al. A phase II clinical trial to analyze olaparib response in patients with BRCA1 and/or BRCA2 promoter methylation with advanced breast cancer (GEICAM/2015-06 COMETA-Breast study). *J Clin Oncol*. 2018;36(15_suppl):TPS1114-TPS1114. doi:10.1200/JCO.2018.36.15_suppl.TPS1114
47. Molinaro A, Becattini B, Mazzoli A, et al. Insulin-Driven PI3K-AKT Signaling in the Hepatocyte Is Mediated by Redundant PI3K α and PI3K β Activities and Is Promoted by RAS. *Cell Metab*. 2019;29(6):1400-1409.e5. doi:10.1016/j.cmet.2019.03.010
48. Yuan Y, Wen W, Yost SE, et al. Combination therapy with BYL719 and LEE011 is synergistic and causes a greater suppression of p-S6 in triple negative breast cancer. *Sci Rep*. 2019;9(1):7509-7509. doi:10.1038/s41598-019-43429-7
49. Kriegsmann M, Endris V, Wolf T, et al. Mutational profiles in triple-negative breast cancer defined by ultradeep multigene sequencing show high rates of PI3K pathway alterations and clinically relevant entity subgroup specific differences. *Oncotarget*. 2014;5(20):9952-9965. doi:10.18632/oncotarget.2481
50. Cossu-Rocca P, Orrù S, Muroi MR, et al. Analysis of PIK3CA Mutations and Activation Pathways in Triple Negative Breast Cancer. Phillips WA, ed. *PLOS ONE*. 2015;10(11):e0141763-e0141763. doi:10.1371/journal.pone.0141763
51. Shah SP, Roth A, Goya R, et al. The clonal and mutational evolution spectrum of primary triple-negative breast cancers. *Nature*. 2012;486(7403):395-399. doi:10.1038/nature10933
52. Carpenter CL, Duckworth BC, Auger KR, Cohen B, Schaffhausen BS, Cantley LC. Purification and characterization of phosphoinositide 3-kinase from rat liver. *J Biol Chem*. 1990;265(32):19704-19711. <http://www.ncbi.nlm.nih.gov/pubmed/2174051>
53. Engelman JA, Luo J, Cantley LC. The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. *Nat Rev Genet*. 2006;7(8):606-619. doi:10.1038/nrg1879

54. Janku F, Yap TA, Meric-Bernstam F. Targeting the PI3K pathway in cancer: are we making headway? *Nat Rev Clin Oncol*. 2018;15(5):273-291. doi:10.1038/nrclinonc.2018.28
55. Maehama T, Dixon JE. The tumor suppressor, PTEN/MMAC1, dephosphorylates the lipid second messenger, phosphatidylinositol 3,4,5-trisphosphate. *J Biol Chem*. 1998;273(22):13375-13378. doi:10.1074/jbc.273.22.13375
56. Stambolic V, Suzuki A, de la Pompa JL, et al. Negative regulation of PKB/Akt-dependent cell survival by the tumor suppressor PTEN. *Cell*. 1998;95(1):29-39. doi:10.1016/s0092-8674(00)81780-8
57. Sun H, Lesche R, Li DM, et al. PTEN modulates cell cycle progression and cell survival by regulating phosphatidylinositol 3,4,5-trisphosphate and Akt/protein kinase B signaling pathway. *Proc Natl Acad Sci U S A*. 1999;96(11):6199-6204. doi:10.1073/pnas.96.11.6199
58. Li J, Yen C, Liaw D, et al. PTEN, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. *Science*. 1997;275(5308):1943-1947. doi:10.1126/science.275.5308.1943
59. García JM, Silva J, Peña C, et al. Promoter methylation of the PTEN gene is a common molecular change in breast cancer. *Genes Chromosomes Cancer*. 2004;41(2):117-124. doi:10.1002/gcc.20062
60. Fritsch C, Huang A, Chatenay-Rivauday C, et al. Characterization of the novel and specific PI3K α inhibitor NVP-BYL719 and development of the patient stratification strategy for clinical trials. *Mol Cancer Ther*. 2014;13(5):1117-1129. doi:10.1158/1535-7163.MCT-13-0865
61. Tan X, Zhang Z, Liu P, Yao H, tong jingshan. *Alpelisib Inhibits PIK3CA-Mutant Breast Cancer Growth Through AKT-Dependent Bim Induction and Mcl-1 Degradation*. In Review; 2020. doi:10.21203/rs.3.rs-38718/v1
62. U.S. Food & Drug Administration. FDA approves alpelisib for metastatic breast cancer. Published 2019. <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-alpelisib-metastatic-breast-cancer>
63. European Medicines Agency. Piqray. Published 2020. <https://www.ema.europa.eu/en/medicines/human/EPAR/piqray>
64. Martín M, Chan A, Dirix L, et al. A randomized adaptive phase II/III study of buparlisib, a pan-class I PI3K inhibitor, combined with paclitaxel for the treatment of HER2– advanced breast cancer (BELLE-4). *Ann Oncol*. 2017;28(2):313-320. doi:10.1093/annonc/mdw562
65. Pascual J, Turner NC. Targeting the PI3-kinase pathway in triple-negative breast cancer. *Ann Oncol*. 2019;30(7):1051-1060. doi:10.1093/annonc/mdz133
66. Juvekar A, Burga LN, Hu H, et al. Combining a PI3K Inhibitor with a PARP Inhibitor Provides an Effective Therapy for BRCA1-Related Breast Cancer. *Cancer Discov*. 2012;2(11):1048-1063. doi:10.1158/2159-8290.CD-11-0336

67. Ibrahim YH, García-García C, Serra V, et al. PI3K Inhibition Impairs BRCA1/2 Expression and Sensitizes BRCA-Proficient Triple-Negative Breast Cancer to PARP Inhibition. *Cancer Discov.* 2012;2(11):1036-1047. doi:10.1158/2159-8290.CD-11-0348
68. Juvekar A, Hu H, Yadegarynia S, et al. Phosphoinositide 3-kinase inhibitors induce DNA damage through nucleoside depletion. *Proc Natl Acad Sci.* 2016;113(30):E4338-E4347. doi:10.1073/pnas.1522223113
69. Konstantinopoulos PA, Barry WT, Birrer M, et al. Olaparib and α -specific PI3K inhibitor alpelisib for patients with epithelial ovarian cancer: a dose-escalation and dose-expansion phase 1b trial. *Lancet Oncol.* 2019;20(4):570-580. doi:10.1016/S1470-2045(18)30905-7
70. Hsu H, Lacey DL, Dunstan CR, et al. Tumor necrosis factor receptor family member RANK mediates osteoclast differentiation and activation induced by osteoprotegerin ligand. *Proc Natl Acad Sci U S A.* 1999;96(7):3540-3545. doi:10.1073/pnas.96.7.3540
71. Lacey DL, Timms E, Tan HL, et al. Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell.* 1998;93(2):165-176. doi:10.1016/s0092-8674(00)81569-x
72. Wong BR, Josien R, Lee SY, et al. TRANCE (tumor necrosis factor [TNF]-related activation-induced cytokine), a new TNF family member predominantly expressed in T cells, is a dendritic cell-specific survival factor. *J Exp Med.* 1997;186(12):2075-2080. doi:10.1084/jem.186.12.2075
73. Yasuda H, Shima N, Nakagawa N, et al. Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. *Proc Natl Acad Sci U S A.* 1998;95(7):3597-3602. doi:10.1073/pnas.95.7.3597
74. Anderson DM, Maraskovsky E, Billingsley WL, et al. A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function. *Nature.* 1997;390(6656):175-179. doi:10.1038/36593
75. Ikeda T, Kasai M, Utsuyama M, Hirokawa K. Determination of Three Isoforms of the Receptor Activator of Nuclear Factor- κ B Ligand and Their Differential Expression in Bone and Thymus 1. *Endocrinology.* 2001;142(4):1419-1426. doi:10.1210/endo.142.4.8070
76. Lum L, Wong BR, Josien R, et al. Evidence for a Role of a Tumor Necrosis Factor- α (TNF- α)-converting Enzyme-like Protease in Shedding of TRANCE, a TNF Family Member Involved in Osteoclastogenesis and Dendritic Cell Survival. *J Biol Chem.* 1999;274(19):13613-13618. doi:10.1074/jbc.274.19.13613
77. Hikita A, Yana I, Wakeyama H, et al. Negative Regulation of Osteoclastogenesis by Ectodomain Shedding of Receptor Activator of NF- κ B Ligand. *J Biol Chem.* 2006;281(48):36846-36855. doi:10.1074/jbc.M606656200
78. Lynch CC, Hikosaka A, Acuff HB, et al. MMP-7 promotes prostate cancer-induced osteolysis via the solubilization of RANKL. *Cancer Cell.* 2005;7(5):485-496. doi:10.1016/j.ccr.2005.04.013
79. Kartsogiannis V, Zhou H, Horwood NJ, et al. Localization of RANKL (receptor activator of NF κ B ligand) mRNA and protein in skeletal and extraskeletal tissues. *Bone.* 1999;25(5):525-534. doi:10.1016/S8756-3282(99)00214-8

80. Collin-Osdoby P, Rothe L, Anderson F, Nelson M, Maloney W, Osdoby P. Receptor Activator of NF- κ B and Osteoprotegerin Expression by Human Microvascular Endothelial Cells, Regulation by Inflammatory Cytokines, and Role in Human Osteoclastogenesis. *J Biol Chem.* 2001;276(23):20659-20672. doi:10.1074/jbc.M010153200
81. Fata JE, Kong Y-Y, Li J, et al. *The Osteoclast Differentiation Factor Osteoprotegerin-Ligand Is Essential for Mammary Gland Development.* Vol 103.; 2000:41-50. doi:10.1016/s0092-8674(00)00103-3
82. Ming J, Cronin SJF, Penninger JM. Targeting the RANKL/RANK/OPG Axis for Cancer Therapy. *Front Oncol.* 2020;10:1283. doi:10.3389/fonc.2020.01283
83. Theoleyre S, Wittrant Y, Tat SK, Fortun Y, Redini F, Heymann D. The molecular triad OPG/RANK/RANKL: involvement in the orchestration of pathophysiological bone remodeling. *Cytokine Growth Factor Rev.* 2004;15(6):457-475. doi:10.1016/j.cytogfr.2004.06.004
84. Wong BR, Josien R, Lee SY, Vologodskaya M, Steinman RM, Choi Y. The TRAF family of signal transducers mediates NF- κ B activation by the TRANCE receptor. *J Biol Chem.* 1998;273(43):28355-28359. doi:10.1074/jbc.273.43.28355
85. Galibert L, Tometsko ME, Anderson DM, Cosman D, Dougall WC. The Involvement of Multiple Tumor Necrosis Factor Receptor (TNFR)-associated Factors in the Signaling Mechanisms of Receptor Activator of NF- κ B, a Member of the TNFR Superfamily. *J Biol Chem.* 1998;273(51):34120-34127. doi:10.1074/jbc.273.51.34120
86. Raju R, Balakrishnan L, Nanjappa V, et al. A comprehensive manually curated reaction map of RANKL/RANK-signaling pathway. *Database J Biol Databases Curation.* 2011;2011:bar021-bar021. doi:10.1093/database/bar021
87. Darnay BG, Ni J, Moore PA, Aggarwal BB. Activation of NF- κ B by RANK requires tumor necrosis factor receptor-associated factor (TRAF) 6 and NF- κ B-inducing kinase. Identification of a novel TRAF6 interaction motif. *J Biol Chem.* 1999;274(12):7724-7731. doi:10.1074/jbc.274.12.7724
88. Darnay BG, Haridas V, Ni J, Moore PA, Aggarwal BB. Characterization of the intracellular domain of receptor activator of NF- κ B (RANK). Interaction with tumor necrosis factor receptor-associated factors and activation of NF- κ B and c-Jun N-terminal kinase. *J Biol Chem.* 1998;273(32):20551-20555. doi:10.1074/jbc.273.32.20551
89. Wong BR, Besser D, Kim N, et al. TRANCE, a TNF Family Member, Activates Akt/PKB through a Signaling Complex Involving TRAF6 and c-Src. *Mol Cell.* 1999;4(6):1041-1049. doi:10.1016/S1097-2765(00)80232-4
90. Papanastasiou AD, Sirinian C, Kalofonos HP. Identification of novel human receptor activator of nuclear factor- κ B isoforms generated through alternative splicing: implications in breast cancer cell survival and migration. *Breast Cancer Res.* 2012;14(4):R112-R112. doi:10.1186/bcr3234
91. Sirinian C, Papanastasiou AD, Schizas M, et al. RANK-c attenuates aggressive properties of ER-negative breast cancer by inhibiting NF- κ B activation and EGFR signaling. *Oncogene.* 2018;37(37):5101-5114. doi:10.1038/s41388-018-0324-y

92. Simonet WS, Lacey DL, Dunstan CR, et al. Osteoprotegerin: a novel secreted protein involved in the regulation of bone density. *Cell*. 1997;89(2):309-319. doi:10.1016/s0092-8674(00)80209-3
93. Baud'huin M, Duplomb L, Téletchéa S, et al. Factor VIII-von Willebrand Factor Complex Inhibits Osteoclastogenesis and Controls Cell Survival. *J Biol Chem*. 2009;284(46):31704-31713. doi:10.1074/jbc.M109.030312
94. Tat SK, Padrines M, Théoleyre S, et al. OPG/membranous-RANKL complex is internalized via the clathrin pathway before a lysosomal and a proteasomal degradation. *Bone*. 2006;39(4):706-715. doi:10.1016/j.bone.2006.03.016
95. Nelson CA, Warren JT, Wang MW-H, Teitelbaum SL, Fremont DH. RANKL Employs Distinct Binding Modes to Engage RANK and the Osteoprotegerin Decoy Receptor. *Structure*. 2012;20(11):1971-1982. doi:10.1016/J.STR.2012.08.030
96. Yasuda H, Shima N, Nakagawa N, et al. Identity of Osteoclastogenesis Inhibitory Factor (OCIF) and Osteoprotegerin (OPG): A Mechanism by which OPG/OCIF Inhibits Osteoclastogenesis *in Vitro*¹. *Endocrinology*. 1998;139(3):1329-1337. doi:10.1210/endo.139.3.5837
97. Holen I, Croucher PI, Hamdy FC, Eaton CL. Osteoprotegerin (OPG) is a survival factor for human prostate cancer cells. *Cancer Res*. 2002;62(6):1619-1623.
98. Théoleyre S, Kwan Tat S, Vusio P, et al. Characterization of osteoprotegerin binding to glycosaminoglycans by surface plasmon resonance: Role in the interactions with receptor activator of nuclear factor κ B ligand (RANKL) and RANK. *Biochem Biophys Res Commun*. 2006;347(2):460-467. doi:10.1016/j.bbrc.2006.06.120
99. Luo J, Yang Z, Ma Y, et al. LGR4 is a receptor for RANKL and negatively regulates osteoclast differentiation and bone resorption. *Nat Med*. 2016;22(5). doi:10.1038/nm.4076
100. Dougall WC, Glaccum M, Charrier K, et al. RANK is essential for osteoclast and lymph node development. *Genes Dev*. 1999;13(18):2412-2424. doi:10.1101/gad.13.18.2412
101. Akiyama T, Shimo Y, Yanai H, et al. The Tumor Necrosis Factor Family Receptors RANK and CD40 Cooperatively Establish the Thymic Medullary Microenvironment and Self-Tolerance. *Immunity*. 2008;29(3):423-437. doi:10.1016/J.IMMUNI.2008.06.015
102. Riggs BL, Khosla S, Melton LJ. A unitary model for involutional osteoporosis: estrogen deficiency causes both type I and type II osteoporosis in postmenopausal women and contributes to bone loss in aging men. *J Bone Miner Res Off J Am Soc Bone Miner Res*. 1998;13(5):763-773. doi:10.1359/jbmr.1998.13.5.763
103. Hanada R. The role of the RANKL/RANK/OPG system in the central nervous systems (CNS). *J Bone Miner Metab*. Published online September 4, 2020. doi:10.1007/s00774-020-01143-9
104. Shackleton M, Vaillant F, Simpson KJ, et al. Generation of a functional mammary gland from a single stem cell. *Nature*. 2006;439(7072):84-88. doi:10.1038/nature04372

105. Asselin-Labat M-L, Vaillant F, Shackleton M, Bouras T, Lindeman GJ, Visvader JE. Delineating the Epithelial Hierarchy in the Mouse Mammary Gland. *Cold Spring Harb Symp Quant Biol.* 2008;73(0):469-478. doi:10.1101/sqb.2008.73.020
106. Mesen TB, Young SL. Progesterone and the Luteal Phase. *Obstet Gynecol Clin North Am.* 2015;42(1):135-151. doi:10.1016/j.ogc.2014.10.003
107. Tanos T, Sflomos G, Echeverria PC, et al. Progesterone/RANKL is a major regulatory axis in the human breast. *Sci Transl Med.* 2013;5(182):182ra55-182ra55. doi:10.1126/scitranslmed.3005654
108. Cao Y, Bonizzi G, Seagroves TN, et al. IKK α Provides an Essential Link between RANK Signaling and Cyclin D1 Expression during Mammary Gland Development. *Cell.* 2001;107(6):763-775. doi:10.1016/S0092-8674(01)00599-2
109. Belet M, Rajaram RD, Caikovski M, et al. Two distinct mechanisms underlie progesterone-induced proliferation in the mammary gland. *Proc Natl Acad Sci U S A.* 2010;107(7):2989-2994. doi:10.1073/pnas.0915148107
110. Kim N-S, Kim H-J, Koo B-K, et al. Receptor activator of NF-kappaB ligand regulates the proliferation of mammary epithelial cells via Id2. *Mol Cell Biol.* 2006;26(3):1002-1013. doi:10.1128/MCB.26.3.1002-1013.2006
111. Asselin-Labat M-L, Vaillant F, Sheridan JM, et al. Control of mammary stem cell function by steroid hormone signalling. *Nature.* 2010;465(7299):798-802. doi:10.1038/nature09027
112. Infante M, Fabi A, Cognetti F, Gorini S, Caprio M, Fabbri A. RANKL/RANK/OPG system beyond bone remodeling: involvement in breast cancer and clinical perspectives. *J Exp Clin Cancer Res.* 2019;38(1):12-12. doi:10.1186/s13046-018-1001-2
113. Santini D, Perrone G, Roato I, et al. Expression pattern of receptor activator of NF κ B (RANK) in a series of primary solid tumors and related bone metastases. *J Cell Physiol.* 2011;226(3):780-784. doi:10.1002/jcp.22402
114. Roux S, Meignin V, Quillard J, et al. RANK (receptor activator of nuclear factor- κ B) and RANKL expression in multiple myeloma: RANKL and RANK in Myeloma. *Br J Haematol.* 2002;117(1):86-92. doi:10.1046/j.1365-2141.2002.03417.x
115. Heymann M-F, Riet A, Le Goff B, Battaglia S, Paineau J, Heymann D. OPG, RANK and RANK ligand expression in thyroid lesions. *Regul Pept.* 2008;148(1-3):46-53. doi:10.1016/j.regpep.2008.02.004
116. Reyes ME, Fujii T, Branstetter D, et al. Poor prognosis of patients with triple-negative breast cancer can be stratified by RANK and RANKL dual expression. *Breast Cancer Res Treat.* 2017;164(1):57-67. doi:10.1007/s10549-017-4233-5
117. Chen G, Sircar K, Aprikian A, Potti A, Goltzman D, Rabbani SA. Expression of RANKL/RANK/OPG in primary and metastatic human prostate cancer as markers of disease stage and functional regulation. *Cancer.* 2006;107(2):289-298. doi:10.1002/cncr.21978

118. Zhang X, Song Y, Song N, et al. Rankl expression predicts poor prognosis in gastric cancer patients: results from a retrospective and single-center analysis. *Braz J Med Biol Res.* 2018;51(3):e6265. doi:10.1590/1414-431x20176265
119. Tsubaki M, Komai M, Fujimoto S, et al. Activation of NF- κ B by the RANKL/RANK system up-regulates snail and twist expressions and induces epithelial-to-mesenchymal transition in mammary tumor cell lines. *J Exp Clin Cancer Res.* 2013;32(1):62-62. doi:10.1186/1756-9966-32-62
120. Gomes I, De Almeida BP, Dâmaso S, et al. Expression of receptor activator of NF κ B (RANK) drives stemness and resistance to therapy in ER+HER2-breast cancer. *Oncotarget.* 2020;11(19):1714-1728. www.oncotarget.com
121. Chen L-M, Kuo C-H, Lai T-Y, et al. RANKL increases migration of human lung cancer cells through intercellular adhesion molecule-1 up-regulation. *J Cell Biochem.* 2011;112(3):933-941. doi:10.1002/jcb.23009
122. Li X, Liu Y, Wu B, et al. Potential role of the OPG/RANK/RANKL axis in prostate cancer invasion and bone metastasis. *Oncol Rep.* 2014;32(6):2605-2611. doi:10.3892/or.2014.3511
123. Casimiro S, Mohammad KS, Pires R, et al. RANKL/RANK/MMP-1 Molecular Triad Contributes to the Metastatic Phenotype of Breast and Prostate Cancer Cells In Vitro. *PLoS ONE.* 2013;8(5). doi:10.1371/journal.pone.0063153
124. Fujimura T, Kambayashi Y, Furudate S, Asano M, Kakizaki A, Aiba S. Receptor Activator of NF- κ B Ligand Promotes the Production of CCL17 from RANK+ M2 Macrophages. *J Invest Dermatol.* 2015;135(11):2884-2887. doi:10.1038/jid.2015.209
125. Palafox M, Ferrer I, Pellegrini P, et al. RANK Induces Epithelial-Mesenchymal Transition and Stemness in Human Mammary Epithelial Cells and Promotes Tumorigenesis and Metastasis. *Cancer Res.* 2012;72(11):2879-2888. doi:10.1158/0008-5472.CAN-12-0044
126. Schramek D, Leibbrandt A, Sigl V, et al. Osteoclast differentiation factor RANKL controls development of progestin-driven mammary cancer. *Nature.* 2010;468(7320):98-102. doi:10.1038/nature09387
127. Gonzalez-Suarez E, Jacob AP, Jones J, et al. RANK ligand mediates progestin-induced mammary epithelial proliferation and carcinogenesis. *Nature.* 2010;468(7320):103-107. doi:10.1038/nature09495
128. Pellegrini P, Cordero A, Gallego MI, et al. Constitutive activation of RANK disrupts mammary cell fate leading to tumorigenesis: RANK Disrupts Mammary Cell Fate Leading to Cancer. *STEM CELLS.* 2013;31(9):1954-1965. doi:10.1002/stem.1454
129. Nolan E, Vaillant F, Branstetter D, et al. RANK ligand as a potential target for breast cancer prevention in BRCA1-mutation carriers. *Nat Med.* 2016;22(8):933-939. doi:10.1038/nm.4118
130. Sigl V, Owusu-Boaitey K, Joshi PA, et al. RANKL/RANK control Brcal mutation-driven mammary tumors. *Cell Res.* 2016;26(7):761-774. doi:10.1038/cr.2016.69

131. Rao S, Cronin SJF, Sigl V, Penninger JM. RANKL and RANK: From Mammalian Physiology to Cancer Treatment. *Trends Cell Biol.* 2018;28(3):213-223. doi:10.1016/j.tcb.2017.11.001
132. Blake ML, Tometsko M, Miller R, Jones JC, Dougall WC. RANK expression on breast cancer cells promotes skeletal metastasis. *Clin Exp Metastasis.* 2014;31(2):233-245. doi:10.1007/s10585-013-9624-3
133. Sun X, Cheng G, Hao M, et al. CXCL12 / CXCR4 / CXCR7 chemokine axis and cancer progression. *Cancer Metastasis Rev.* 2010;29(4):709-722. doi:10.1007/s10555-010-9256-x
134. Soki FN, Park SI, McCauley LK. The multifaceted actions of PTHrP in skeletal metastasis. *Future Oncol.* 2012;8(7):803-817. doi:10.2217/fon.12.76
135. Le Pape F, Vargas G, Clézardin P. The role of osteoclasts in breast cancer bone metastasis. *J Bone Oncol.* 2016;5(3):93-95. doi:10.1016/j.jbo.2016.02.008
136. Wu X, Li F, Dang L, Liang C, Lu A, Zhang G. RANKL/RANK System-Based Mechanism for Breast Cancer Bone Metastasis and Related Therapeutic Strategies. *Front Cell Dev Biol.* 2020;8. doi:10.3389/fcell.2020.00076
137. de Groot AF, Appelman-Dijkstra NM, van der Burg SH, Kroep JR. The anti-tumor effect of RANKL inhibition in malignant solid tumors – A systematic review. *Cancer Treat Rev.* 2018;62:18-28. doi:10.1016/j.ctrv.2017.10.010
138. Paget S. The distribution of secondary growths in cancer of the breast. *The Lancet.* 1889;133(3421):571-573. doi:10.1016/S0140-6736(00)49915-0
139. European Medicines Agency. Prolia. Published 2010. <https://www.ema.europa.eu/en/medicines/human/EPAR/prolia>
140. U.S. Food & Drug Administration. Prolia - label. Published 2010. https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/125320s186lbl.pdf
141. U.S. Food & Drug Administration. Xgeva - label. Published 2010. https://www.accessdata.fda.gov/drugsatfda_docs/label/2013/125320s094lbl.pdf
142. European Medicines Agency. Xgeva. Published 2011. <https://www.ema.europa.eu/en/medicines/human/EPAR/xgeva>
143. Scagliotti GV, Hirsh V, Siena S, et al. Overall Survival Improvement in Patients with Lung Cancer and Bone Metastases Treated with Denosumab Versus Zoledronic Acid: Subgroup Analysis from a Randomized Phase 3 Study. *J Thorac Oncol.* 2012;7(12):1823-1829. doi:10.1097/JTO.0b013e31826aec2b
144. Rossini M, Viapiana O, Adami S, et al. Effects of denosumab on peripheral lymphocyte subpopulations. *Endocrine.* 2016;53(3):857-859. doi:10.1007/s12020-015-0723-6
145. Matlock K, Berlow N, Keller C, Pal R. Combination therapy design for maximizing sensitivity and minimizing toxicity. *BMC Bioinformatics.* 2017;18(S4):116. doi:10.1186/s12859-017-1523-1

146. Sun W, Sanderson PE, Zheng W. Drug combination therapy increases successful drug repositioning. *Drug Discov Today*. 2016;21(7):1189-1195. doi:10.1016/j.drudis.2016.05.015
147. Lee K, Chung YH, Ahn H, Kim H, Rho J, Jeong D. Selective Regulation of MAPK Signaling Mediates RANKL-dependent Osteoclast Differentiation. *Int J Biol Sci*. 2016;12(2):235-245. doi:10.7150/ijbs.13814
148. Zauli G, Melloni E, Capitani S, Secchiero P. Role of full-length osteoprotegerin in tumor cell biology. *Cell Mol Life Sci CMLS*. 2009;66(5):841-851. doi:10.1007/s00018-008-8536-x
149. R&D Systems. Human TRANCE/TNFSF11/RANK L Antibody. https://www.rndsystems.com/products/human-trance-tnfsf11-rank-l-antibody-70525_mab626#product-datasheets
150. Keung MY, Wu Y, Badar F, Vadgama JV. Response of Breast Cancer Cells to PARP Inhibitors Is Independent of BRCA Status. *J Clin Med*. 2020;9(4):940. doi:10.3390/jcm9040940
151. Arun B, Akar U, Gutierrez-Barrera AM, Hortobagyi GN, Ozpolat B. The PARP inhibitor AZD2281 (Olaparib) induces autophagy/mitophagy in BRCA1 and BRCA2 mutant breast cancer cells. *Int J Oncol*. 2015;47(1):262-268. doi:10.3892/ijo.2015.3003
152. Noordermeer SM, van Attikum H. PARP Inhibitor Resistance: A Tug-of-War in BRCA-Mutated Cells. *Trends Cell Biol*. 2019;29(10):820-834. doi:10.1016/j.tcb.2019.07.008
153. Juric D, Castel P, Griffith M, et al. Convergent loss of PTEN leads to clinical resistance to a PI(3)K α inhibitor. *Nature*. 2015;518(7538):240-244. doi:10.1038/nature13948
154. Costa C, Ebi H, Martini M, et al. Measurement of PIP3 Levels Reveals an Unexpected Role for p110 β in Early Adaptive Responses to p110 α -Specific Inhibitors in Luminal Breast Cancer. *Cancer Cell*. 2015;27(1):97-108. doi:10.1016/j.ccell.2014.11.007
155. Clement E, Inuzuka H, Nihira NT, Wei W, Toker A. Skp2-dependent reactivation of AKT drives resistance to PI3K inhibitors. *Sci Signal*. 2018;11(521):eaao3810. doi:10.1126/scisignal.aao3810
156. Ahern E, Harjunpää H, O'Donnell JS, et al. RANKL blockade improves efficacy of PD1-PD-L1 blockade or dual PD1-PD-L1 and CTLA4 blockade in mouse models of cancer. *OncImmunology*. 2018;7(6):e1431088. doi:10.1080/2162402X.2018.1431088
157. Luen S, Virassamy B, Savas P, Salgado R, Loi S. The genomic landscape of breast cancer and its interaction with host immunity. *The Breast*. 2016;29:241-250. doi:10.1016/j.breast.2016.07.015
158. Mittendorf EA, Philips AV, Meric-Bernstam F, et al. PD-L1 Expression in Triple-Negative Breast Cancer. *Cancer Immunol Res*. 2014;2(4):361-370. doi:10.1158/2326-6066.CIR-13-0127
159. Denkert C, von Minckwitz G, Darb-Esfahani S, et al. Tumour-infiltrating lymphocytes and prognosis in different subtypes of breast cancer: a pooled analysis of 3771 patients treated with neoadjuvant therapy. *Lancet Oncol*. 2018;19(1):40-50. doi:10.1016/S1470-2045(17)30904-X
160. ATCC. Triple Negative Breast Cancer Panels. <http://www.atcc.org/~media/03E3D0D9F0C74FE593CE7D010F982CA2.ashx>

161. Saal LH, Gruvberger-Saal SK, Persson C, et al. Recurrent gross mutations of the PTEN tumor suppressor gene in breast cancers with deficient DSB repair. *Nat Genet.* 2008;40(1):102-107. doi:10.1038/ng.2007.39
162. The Broad Institute of MIT & Harvard. CCLE (Cancer Cell Line Encyclopedia). Published 2020. <https://portals.broadinstitute.org/ccle>
163. Elkabets M, Vora S, Juric D, et al. mTORC1 inhibition is required for sensitivity to PI3K p110 α inhibitors in PIK3CA-mutant breast cancer. *Sci Transl Med.* 2013;5(196):196ra99-196ra99. doi:10.1126/scitranslmed.3005747
164. Pierce A, McGowan PM, Cotter M, et al. Comparative antiproliferative effects of iniparib and olaparib on a panel of triple-negative and non-triple-negative breast cancer cell lines. *Cancer Biol Ther.* 2013;14(6):537-545. doi:10.4161/cbt.24349
165. O'Brien NA, McDonald K, Tong L, et al. Targeting PI3K/mTOR overcomes resistance to HER2-targeted therapy independent of feedback activation of AKT. *Clin Cancer Res.* 2014;20(13):3507-3520. doi:10.1158/1078-0432.CCR-13-2769
166. Xu Y, Wang X, Chen Y, et al. Integration of Receptor Tyrosine Kinases Determines Sensitivity to PI3K α -selective Inhibitors in Breast Cancer. *Theranostics.* 2017;7(4):974-986. doi:10.7150/thno.17830

APPENDIX

Table S1 *BRCA1/2*, *PTEN* and *PIK3CA* status of the TNBC cell lines in the panel^{160,161}.

Cell line	<i>BRCA1/2</i> status	<i>PTEN</i> status	<i>PIK3CA</i> status
HCC1937	<i>BRCA1</i> mutated (homozygous c.5266_5267insC)	Mutated (homozygous deletion)	Wild type
BT-20	Wild type	Wild type	Mutated (heterozygous c.1616C>G and heterozygous c.3140A>G)
MDA-MB-231	Wild type	Wild type	Wild type
MDA-MB-468	Wild type	Mutated (homozygous c.253+1G>T)	Wild type

Table S2 mRNA expression data for *RANK* (*TNFRSF11A*) and *RANKL* (*TNFSF11*) across BC cell lines, retrieved from the Cancer Cell Line Encyclopedia (CCLE)^{120,162}. Values were obtained via RNASeq and are presented in Log2RPKM ordered for decreasing *RANK* expression. TNBC cells used in this project are highlighted.

Cell line	<i>TNFRSF11A</i>	<i>TNFSF11</i>
HCC70_BREAST	2.9401096	-0.46621224
AU565_BREAST	1.398290561	-13
HCC2157_BREAST	1.331681415	-13
HCC1143_BREAST	1.19790117	-5.106361658
HCC1187_BREAST	1.189016004	-4.856818136
CAL51_BREAST	1.057985114	-13
HCC1954_BREAST	0.973932394	-3.902250589
MDAMB468_BREAST	0.904624149	-13
MDAMB231_BREAST	0.838153185	-13
MDAMB175VII_BREAST	0.491170875	-4.534546938
HCC38_BREAST	0.149388964	-4.103090872
ZR751_BREAST	0.033290241	-13
HMEL_BREAST	-0.056751978	-3.674601071
BT20_BREAST	-0.0593241	-13
CAL851_BREAST	-0.233893939	-4.885490814
HCC1599_BREAST	-0.284225335	-3.534351023
SKBR3_BREAST	-0.420929371	-6.807569465
MDAMB415_BREAST	-0.498624298	-3.37132519
HMC18_BREAST	-0.853397765	-13
HCC1937_BREAST	-0.95089629	-0.796249793
EFM192A_BREAST	-1.258191943	-7.19868407
HCC1428_BREAST	-1.284746289	-13
MDAMB361_BREAST	-1.376981128	-13
HCC1419_BREAST	-1.502619572	-13
KPL1_BREAST	-1.531007627	-13

MCF7_BREAST	-1.557351949	-13
HCC202_BREAST	-1.567411078	-13
DU4475_BREAST	-1.579180223	-13
HCC1806_BREAST	-1.681640483	-5.94547517
HCC2218_BREAST	-1.705042634	-13
T47D_BREAST	-1.793258066	-13
UACC812_BREAST	-1.921447176	-13
HCC1395_BREAST	-1.973981675	-13
MDAMB436_BREAST	-2.104042041	-7.197323212
CAMA1_BREAST	-2.142396515	-13
BT549_BREAST	-2.244616152	-4.902511133
HS578T_BREAST	-2.246154962	-13
BT474_BREAST	-2.382347132	-7.05113046
HDQP1_BREAST	-2.526858607	-0.807856812
CAL120_BREAST	-2.715849158	-13
MDAMB453_BREAST	-2.779087536	-6.829299985
HCC1569_BREAST	-2.896640629	-13
UACC893_BREAST	-3.557200107	-13
JIMT1_BREAST	-3.646321172	-13
HS274T_BREAST	-3.653719341	-13
MDAMB134VI_BREAST	-3.693672779	-13
CAL148_BREAST	-4.245191587	-13
MDAMB157_BREAST	-4.918962961	-13
HS606T_BREAST	-5.098164082	-13
HS343T_BREAST	-5.374366396	-13
HCC1500_BREAST	-5.390897142	-13
HS742T_BREAST	-6.16185884	-13
HS281T_BREAST	-6.174836524	-13
HS739T_BREAST	-6.551758519	-13
BT483_BREAST	-6.691730693	-13
EFM19_BREAST	-7.179649613	-13
ZR7530_BREAST	-7.713275303	-6.134131208

Table S3 Described IC₅₀ values for olaparib and alpelisib across the TNBC cell lines. Source publication is indicated next to each value.

	HCC1937	BT-20	MDA-MB-231	MDA-MB-468
Olaparib	96 μM ¹⁵⁰	7.7 μM ¹⁶³	> 90 μM ²⁹	5 μM ¹⁶⁴
Alpelisib	> 10 μM ¹⁶⁵	1.3 μM ¹⁶⁶	19.29 μM ⁴⁸	11.28 μM ⁴⁸